

GEORGIAN MEDICAL NEWS

ISSN 1512-0112

NO 3 (348) March 2024

ТБИЛИСИ - NEW YORK



ЕЖЕМЕСЯЧНЫЙ НАУЧНЫЙ ЖУРНАЛ

Медицинские новости Грузии
საქართველოს სამედიცინო სიახლენი

GEORGIAN MEDICAL NEWS

Monthly Georgia-US joint scientific journal published both in electronic and paper formats of the Agency of Medical Information of the Georgian Association of Business Press.
Published since 1994. Distributed in NIS, EU and USA.

GMN: Georgian Medical News is peer-reviewed, published monthly journal committed to promoting the science and art of medicine and the betterment of public health, published by the GMN Editorial Board since 1994. GMN carries original scientific articles on medicine, biology and pharmacy, which are of experimental, theoretical and practical character; publishes original research, reviews, commentaries, editorials, essays, medical news, and correspondence in English and Russian.

GMN is indexed in MEDLINE, SCOPUS, PubMed and VINITI Russian Academy of Sciences. The full text content is available through EBSCO databases.

GMN: Медицинские новости Грузии - ежемесячный рецензируемый научный журнал, издаётся Редакционной коллегией с 1994 года на русском и английском языках в целях поддержки медицинской науки и улучшения здравоохранения. В журнале публикуются оригинальные научные статьи в области медицины, биологии и фармации, статьи обзорного характера, научные сообщения, новости медицины и здравоохранения. Журнал индексируется в MEDLINE, отражён в базе данных SCOPUS, PubMed и ВИНТИ РАН. Полнотекстовые статьи журнала доступны через БД EBSCO.

GMN: Georgian Medical News – საქართველოს სამედიცინო სიახლენი – არის ყოველთვიური სამეცნიერო სამედიცინო რეცენზირებადი ჟურნალი, გამოიცემა 1994 წლიდან, წარმოადგენს სარედაქციო კოლეგიისა და აშშ-ის მეცნიერების, განათლების, ინდუსტრიის, ხელოვნებისა და ბუნებისმეტყველების საერთაშორისო აკადემიის ერთობლივ გამოცემას. GMN-ში რუსულ და ინგლისურ ენებზე ქვეყნდება ექსპერიმენტული, თეორიული და პრაქტიკული ხასიათის ორიგინალური სამეცნიერო სტატიები მედიცინის, ბიოლოგიისა და ფარმაციის სფეროში, მიმოხილვითი ხასიათის სტატიები.

ჟურნალი ინდექსირებულია MEDLINE-ის საერთაშორისო სისტემაში, ასახულია SCOPUS-ის, PubMed-ის და ВИНТИ РАН-ის მონაცემთა ბაზებში. სტატიების სრული ტექსტი ხელმისაწვდომია EBSCO-ს მონაცემთა ბაზებიდან.

WEBSITE

www.geomednews.com

К СВЕДЕНИЮ АВТОРОВ!

При направлении статьи в редакцию необходимо соблюдать следующие правила:

1. Статья должна быть представлена в двух экземплярах, на русском или английском языках, напечатанная через **полтора интервала на одной стороне стандартного листа с шириной левого поля в три сантиметра**. Используемый компьютерный шрифт для текста на русском и английском языках - **Times New Roman (Кириллица)**, для текста на грузинском языке следует использовать **AcadNusx**. Размер шрифта - **12**. К рукописи, напечатанной на компьютере, должен быть приложен CD со статьей.

2. Размер статьи должен быть не менее десяти и не более двадцати страниц машинописи, включая указатель литературы и резюме на английском, русском и грузинском языках.

3. В статье должны быть освещены актуальность данного материала, методы и результаты исследования и их обсуждение.

При представлении в печать научных экспериментальных работ авторы должны указывать вид и количество экспериментальных животных, применявшиеся методы обезболивания и усыпления (в ходе острых опытов).

4. К статье должны быть приложены краткое (на полстраницы) резюме на английском, русском и грузинском языках (включающее следующие разделы: цель исследования, материал и методы, результаты и заключение) и список ключевых слов (key words).

5. Таблицы необходимо представлять в печатной форме. Фотокопии не принимаются. **Все цифровые, итоговые и процентные данные в таблицах должны соответствовать таковым в тексте статьи**. Таблицы и графики должны быть озаглавлены.

6. Фотографии должны быть контрастными, фотокопии с рентгенограмм - в позитивном изображении. Рисунки, чертежи и диаграммы следует озаглавить, пронумеровать и вставить в соответствующее место текста **в tiff формате**.

В подписях к микрофотографиям следует указывать степень увеличения через окуляр или объектив и метод окраски или импрегнации срезов.

7. Фамилии отечественных авторов приводятся в оригинальной транскрипции.

8. При оформлении и направлении статей в журнал МНГ просим авторов соблюдать правила, изложенные в «Единых требованиях к рукописям, представляемым в биомедицинские журналы», принятых Международным комитетом редакторов медицинских журналов - <http://www.spinesurgery.ru/files/publish.pdf> и http://www.nlm.nih.gov/bsd/uniform_requirements.html В конце каждой оригинальной статьи приводится библиографический список. В список литературы включаются все материалы, на которые имеются ссылки в тексте. Список составляется в алфавитном порядке и нумеруется. Литературный источник приводится на языке оригинала. В списке литературы сначала приводятся работы, написанные знаками грузинского алфавита, затем кириллицей и латиницей. Ссылки на цитируемые работы в тексте статьи даются в квадратных скобках в виде номера, соответствующего номеру данной работы в списке литературы. Большинство цитированных источников должны быть за последние 5-7 лет.

9. Для получения права на публикацию статья должна иметь от руководителя работы или учреждения визу и сопроводительное отношение, написанные или напечатанные на бланке и заверенные подписью и печатью.

10. В конце статьи должны быть подписи всех авторов, полностью приведены их фамилии, имена и отчества, указаны служебный и домашний номера телефонов и адреса или иные координаты. Количество авторов (соавторов) не должно превышать пяти человек.

11. Редакция оставляет за собой право сокращать и исправлять статьи. Корректур авторам не высылаются, вся работа и сверка проводится по авторскому оригиналу.

12. Недопустимо направление в редакцию работ, представленных к печати в иных издательствах или опубликованных в других изданиях.

При нарушении указанных правил статьи не рассматриваются.

REQUIREMENTS

Please note, materials submitted to the Editorial Office Staff are supposed to meet the following requirements:

1. Articles must be provided with a double copy, in English or Russian languages and typed or computer-printed on a single side of standard typing paper, with the left margin of 3 centimeters width, and 1.5 spacing between the lines, typeface - **Times New Roman (Cyrillic)**, print size - 12 (referring to Georgian and Russian materials). With computer-printed texts please enclose a CD carrying the same file titled with Latin symbols.

2. Size of the article, including index and resume in English, Russian and Georgian languages must be at least 10 pages and not exceed the limit of 20 pages of typed or computer-printed text.

3. Submitted material must include a coverage of a topical subject, research methods, results, and review.

Authors of the scientific-research works must indicate the number of experimental biological species drawn in, list the employed methods of anesthetization and soporific means used during acute tests.

4. Articles must have a short (half page) abstract in English, Russian and Georgian (including the following sections: aim of study, material and methods, results and conclusions) and a list of key words.

5. Tables must be presented in an original typed or computer-printed form, instead of a photocopied version. **Numbers, totals, percentile data on the tables must coincide with those in the texts of the articles.** Tables and graphs must be headed.

6. Photographs are required to be contrasted and must be submitted with doubles. Please number each photograph with a pencil on its back, indicate author's name, title of the article (short version), and mark out its top and bottom parts. Drawings must be accurate, drafts and diagrams drawn in Indian ink (or black ink). Photocopies of the X-ray photographs must be presented in a positive image in **tiff format**.

Accurately numbered subtitles for each illustration must be listed on a separate sheet of paper. In the subtitles for the microphotographs please indicate the ocular and objective lens magnification power, method of coloring or impregnation of the microscopic sections (preparations).

7. Please indicate last names, first and middle initials of the native authors, present names and initials of the foreign authors in the transcription of the original language, enclose in parenthesis corresponding number under which the author is listed in the reference materials.

8. Please follow guidance offered to authors by The International Committee of Medical Journal Editors guidance in its Uniform Requirements for Manuscripts Submitted to Biomedical Journals publication available online at: http://www.nlm.nih.gov/bsd/uniform_requirements.html
http://www.icmje.org/urm_full.pdf

In GMN style for each work cited in the text, a bibliographic reference is given, and this is located at the end of the article under the title "References". All references cited in the text must be listed. The list of references should be arranged alphabetically and then numbered. References are numbered in the text [numbers in square brackets] and in the reference list and numbers are repeated throughout the text as needed. The bibliographic description is given in the language of publication (citations in Georgian script are followed by Cyrillic and Latin).

9. To obtain the rights of publication articles must be accompanied by a visa from the project instructor or the establishment, where the work has been performed, and a reference letter, both written or typed on a special signed form, certified by a stamp or a seal.

10. Articles must be signed by all of the authors at the end, and they must be provided with a list of full names, office and home phone numbers and addresses or other non-office locations where the authors could be reached. The number of the authors (co-authors) must not exceed the limit of 5 people.

11. Editorial Staff reserves the rights to cut down in size and correct the articles. Proof-sheets are not sent out to the authors. The entire editorial and collation work is performed according to the author's original text.

12. Sending in the works that have already been assigned to the press by other Editorial Staffs or have been printed by other publishers is not permissible.

**Articles that Fail to Meet the Aforementioned
Requirements are not Assigned to be Reviewed.**

ავტორთა საქურაღებოლ!

რედაქციაში სტატიის წარმოდგენისას საჭიროა დაიცვათ შემდეგი წესები:

1. სტატია უნდა წარმოადგინოთ 2 ცალად, რუსულ ან ინგლისურ ენებზე დაბეჭდილი სტანდარტული ფურცლის 1 გვერდზე, 3 სმ სიგანის მარცხენა ველისა და სტრიქონებს შორის 1,5 ინტერვალის დაცვით. გამოყენებული კომპიუტერული შრიფტი რუსულ და ინგლისურენოვან ტექსტებში - **Times New Roman (Кириллица)**, ხოლო ქართულენოვან ტექსტში საჭიროა გამოვიყენოთ **AcadNusx**. შრიფტის ზომა – 12. სტატიას თან უნდა ახლდეს CD სტატიით.

2. სტატიის მოცულობა არ უნდა შეადგენდეს 10 გვერდზე ნაკლებს და 20 გვერდზე მეტს ლიტერატურის სიის და რეზიუმეების (ინგლისურ, რუსულ და ქართულ ენებზე) ჩათვლით.

3. სტატიაში საჭიროა გაშუქდეს: საკითხის აქტუალობა; კვლევის მიზანი; საკვლევი მასალა და გამოყენებული მეთოდები; მიღებული შედეგები და მათი განსჯა. ექსპერიმენტული ხასიათის სტატიების წარმოდგენისას ავტორებმა უნდა მიუთითონ საექსპერიმენტო ცხოველების სახეობა და რაოდენობა; გაუტკივარებისა და დაძინების მეთოდები (მწვავე ცდების პირობებში).

4. სტატიას თან უნდა ახლდეს რეზიუმე ინგლისურ, რუსულ და ქართულ ენებზე არანაკლებ ნახევარი გვერდის მოცულობისა (სათაურის, ავტორების, დაწესებულების მითითებით და უნდა შეიცავდეს შემდეგ განყოფილებებს: მიზანი, მასალა და მეთოდები, შედეგები და დასკვნები; ტექსტუალური ნაწილი არ უნდა იყოს 15 სტრიქონზე ნაკლები) და საკვანძო სიტყვების ჩამონათვალი (key words).

5. ცხრილები საჭიროა წარმოადგინოთ ნაბეჭდი სახით. ყველა ციფრული, შემაჯამებელი და პროცენტული მონაცემები უნდა შეესაბამებოდეს ტექსტში მოყვანილს.

6. ფოტოსურათები უნდა იყოს კონტრასტული; სურათები, ნახაზები, დიაგრამები - დასათაურებული, დანომრილი და სათანადო ადგილას ჩასმული. რენტგენოგრამების ფოტოასლები წარმოადგინეთ პოზიტიური გამოსახულებით **tiff** ფორმატში. მიკროფოტოსურათების წარწერებში საჭიროა მიუთითოთ ოკულარის ან ობიექტივის საშუალებით გადიდების ხარისხი, ანათალების შედეგის ან იმპრეგნაციის მეთოდი და აღნიშნოთ სურათის ზედა და ქვედა ნაწილები.

7. სამამულო ავტორების გვარები სტატიაში აღინიშნება ინიციალების თანდართვით, უცხოურისა – უცხოური ტრანსკრიპციით.

8. სტატიას თან უნდა ახლდეს ავტორის მიერ გამოყენებული სამამულო და უცხოური შრომების ბიბლიოგრაფიული სია (ბოლო 5-8 წლის სიღრმით). ანბანური წყობით წარმოდგენილ ბიბლიოგრაფიულ სიაში მიუთითეთ ჯერ სამამულო, შემდეგ უცხოელი ავტორები (გვარი, ინიციალები, სტატიის სათაური, ჟურნალის დასახელება, გამოცემის ადგილი, წელი, ჟურნალის №, პირველი და ბოლო გვერდები). მონოგრაფიის შემთხვევაში მიუთითეთ გამოცემის წელი, ადგილი და გვერდების საერთო რაოდენობა. ტექსტში კვადრატულ ფხიხლებში უნდა მიუთითოთ ავტორის შესაბამისი N ლიტერატურის სიის მიხედვით. მიზანშეწონილია, რომ ციტირებული წყაროების უმეტესი ნაწილი იყოს 5-6 წლის სიღრმის.

9. სტატიას თან უნდა ახლდეს: ა) დაწესებულების ან სამეცნიერო ხელმძღვანელის წარდგინება, დამოწმებული ხელმოწერითა და ბეჭდით; ბ) დარგის სპეციალისტის დამოწმებული რეცენზია, რომელშიც მითითებული იქნება საკითხის აქტუალობა, მასალის საკმაობა, მეთოდის სანდოობა, შედეგების სამეცნიერო-პრაქტიკული მნიშვნელობა.

10. სტატიის ბოლოს საჭიროა ყველა ავტორის ხელმოწერა, რომელთა რაოდენობა არ უნდა აღემატებოდეს 5-ს.

11. რედაქცია იტოვებს უფლებას შეასწოროს სტატია. ტექსტზე მუშაობა და შეჯერება ხდება საავტორო ორიგინალის მიხედვით.

12. დაუშვებელია რედაქციაში ისეთი სტატიის წარდგენა, რომელიც დასაბეჭდად წარდგენილი იყო სხვა რედაქციაში ან გამოქვეყნებული იყო სხვა გამოცემებში.

აღნიშნული წესების დარღვევის შემთხვევაში სტატიები არ განიხილება.

Содержание:

Alla Kyrychenko, Nataliya Tomakh, Vasyl Kornatsky, Olena Lysunets, Oksana Sirenko, Olexandr Kuryata. ACUTE MYOCARDITIS IN YOUNG AGE MIMICKING AS ST-ELEVATION MYOCARDIAL INFARCTION: CASE REPORT.....	6-9
Nikolaos Geropoulos, Polychronis Voultzos, Miltiadis Geropoulos, Fani Tsolaki, Georgios Tagarakis. CENTRALIZATION AND CORRUPTION IN HEALTH PROCUREMENT OF THE SOUTHERN EUROPEAN UNION COUNTRIES.....	10-21
Yerlan Bazargaliyev, Bibigul Tleumagamabetova, Khatimya Kudabayeva, Raikul Kosmuratova. ANALYSIS OF ANTIDIABETIC THERAPY FOR TYPE 2 DIABETES IN PRIMARY HEALTH CARE (WESTERN KAZAKHSTAN).....	22-27
Christina Mary P Paul, Shashikala Manjunatha, Archana Lakshmi PA, Girisha Sharma. A STUDY ON THE INFORMATION TRANSFER AND LONG-TERM PSYCHOLOGICAL IMPACT OF CHILD SEXUAL ABUSE....	28-31
Nino Chomakhashvili, Nino Chikhladze, Nato Pitskhelauri. ERGONOMIC PRACTICE IN DENTAL CLINICS AND MUSCULOSKELETAL DISORDERS AMONG DENTISTS IN GEORGIA.....	32-35
Chnar S. Maarof, Ali S. Dauod, Rachel E. Dunham. PREVALENCE OF PRETERM DELIVERY AMONG WOMEN WHO RECEIVE PROGESTERONE SUPPLEMENTATION DURING PREGNANCY: CROSS-SECTIONAL OBSERVATIONAL STUDY.....	36-39
S.K. Tukeshov, T.A. Baysekeev, E. D. Choi, G.A. Kulushova, M.I. Nazir, N.B. Jaxymbayev, A.A. Turkmenov. OSTEOSYNTHESIS OF COMPLEX COMMUNUTED HAND BONE FRACTURES BY APPLYING THE LACING METHOD (A CLINICAL CASE STUDY)	40-43
Majed A Mohammad, Firas A Jassim, Ali Malik Tiryag. RETROGRADE INTRARENAL LITHOTRIPSY USING DISPOSABLE FLEXIBLE URETEROSCOPE.....	44-46
Olga Samara, Mykhailo Zhylin, Viktoriia Mendelo, Artur Akopian, Nina Bakuridze. THE ROLE OF EMOTIONAL INTELLIGENCE IN THE DIAGNOSIS AND PSYCHOTHERAPY OF MENTAL DISORDERS: AN ANALYSIS OF PRACTICAL APPROACHES.....	47-53
Arnab Sain, Ralph Keita, Arunava Ray, Nauman Manzoor, Arsany Metry, Ahmed Elkilany, Kanishka Wattage, Michele Halasa, Jack Song Chia, Fahad Hussain, Odiamehi Aisabokhale, Zain Sohail, Vivek Deshmukh, Adhish Avasthi. SAFE USE OF INTRA-OPERATIVE TOURNIQUETS IN A DISTRICT HOSPITAL IN THE UK-AN AUDIT STUDY IN ORTHOPAEDIC THEATRES AND REVIEW OF CURRENT LITERATURE.....	54-56
Takuma Hayashi, Ikuo Konishi. POST-COVID-19 INFLAMMATORY RHEUMATOID ARTHRITIS REMISSION.....	57-59
Athraa Essa Ahmed. KNOWLEDGE OF SECONDARY SCHOOL STUDENTS REGARDING PREVENTIVE MEASURES FOR RESPIRATORY INFECTIOUS DISEASE IN TIKRIT CITY.....	60-62
Irakli Gogokhia, Merab Kiladze, Tamar Gogichaishvili, Koba Sakhechidze. FEASIBILITY AND EFFECTIVENESS OF GENERAL ANESTHESIA WITH OPIOIDS VERSUS OPIOID-FREE ANESTHESIA PLUS TRANSVERSUS ABDOMINIS PLANE BLOCK ON POSTOPERATIVE OUTCOMES AFTER MINI GASTRIC BYPASS SURGERY.....	63-71
Anton I. Korbut, Vyacheslav V. Romanov, Vadim V. Klimontov. URINARY EXCRETION OF ALPHA-ACTININ-4 AND TIGHT JUNCTION PROTEIN 1 IN PATIENTS WITH TYPE 2 DIABETES AND DIFFERENT PATTERNS OF CHRONIC KIDNEY DISEASE.....	72-77
Rishu Bansal, Maia Zhamutashvili, Tinatin Gognadze, Natia Jojua, Ekaterine Dolmazishvili. ENTEROHEMORRHAGIC ESCHERICHIA COLI LEADING TO HAEMOLYTIC UREMIC SYNDROME - CASE STUDY AND REVIEW.....	78-80
Ayah J. Mohammed, Entedhar R. Sarhat. PARTIAL PURIFICATION OF GLUTATHIONE PEROXIDASE ENZYME FROM WOMEN WITH BREAST CANCER.....	81-86
Mariam Kekenadze, Nana kvirkvelia, Maia Beridze, Shorena Vashadze. SEROTONIN AND AMYOTROPHIC LATERAL SCLEROSIS (ALS).....	87-90
Arnab Sain, Zain Sohail, Nauman Manzoor, Amir Varasteh, Vivek Deshmukh, Arsany Metry, Fahad Hussain , Ahmed Elkilany, Kanishka Wattage, Michelle Halasa, Jack Chai Song, Ralph Keita, Odiamehi Aisabokhale, Koushik Ghosh. IMPORTANCE OF JOINT LINE RESTORATION IN TOTAL KNEE ARTHROPLASTY.....	91-93
Lurin I, Gorobeiko M, Lovin A, Gorobeyko B, Lovina N, Dinets A. APPLICATION OF ARTIFICIAL INTELLIGENCE IN CIVIL AND MILITARY MEDICINE.....	94-98
Kassim SA Al Neaimy, Okba N Alsarraf, Maes MK Alkhyatt. COMPARATIVE STUDY OF OXIDATIVE STRESS IN PATIENTS WITH B -THALASSEMIA MAJOR ON DEFERASIROX VERSUS DEFEROXAMINETHERAPY.....	99-102

Hinpetch Daungsupawong, Viroj Wiwanitkit. COMMENT ON "A CROSS-SECTIONAL STUDY ON COVID-19 VACCINATION HESITATION AMONG UNIVERSITY STUDENTS."	103-104
Taisa P. Skrypnikova, Petro M. Skrypnykov, Olga V. Gancho, Galina A. Loban', Julia V. Tymoshenko, Vira I. Fedorchenko, Olena A. Pysarenko, Kseniia A. Lazareva, Tetyana A. Khmil, Olga O. Kulai. IMPROVEMENT OF THE METHODOLOGY OF BIOMATERIAL COLLECTION FOR THE DIAGNOSIS OF THE ORAL CAVITY MUCOSADISEASES.	105-108
Mkrtchyan S, Shukuryan A, Dunamalyan R, Sakanyan G, Galstyan H, Chichoyan N, Mardiyan M. CLINICAL SIGNIFICANCE OF CHANGES IN QUALITY OF LIFE INDICATORS AS A METHOD FOR ASSESSING THE EFFECTIVENESS OF ENT HERBAL REMEDIES.	109-116
OSAMA ARIM, Ali Alshalcy, Mohammed Z. Shakir, Omar KO. Agha, Hayder Alhamdany. TRANSPEDICULAR SCREW FIXATION IN DEGENERATIVE LUMBOSACRAL SPINE DISEASE SURGICAL OUTCOME.	117-121
Tavartkiladze G, Kalandadze M, Puturidze S, Parulava Sh, Margvelashvili V. TEMPOROMANDIBULAR JOINT DISORDERS AND THE WAY OF THEIR OPTIMIZATION: A LITERATURE REVIEW.	22-127
Mohammed Saarti, Mohammed D Mahmood, Loay A. Alchalaby. OVERVIEW OF DRUG-INDUCED OROFACIAL CLEFT.	128-131
Tchernev G, Broshtilova V. (NDMA) METFORMIN AND (NTP) SITAGLIPTIN INDUCED CUTANEOUS MELANOMAS: LINKS TO NITROSOGENESIS, NITROSO-PHOTOCARCINOGENESIS, ONCOPHARMACOGENESIS AND THE METABOLIC REPROGRAMMING.	132-143
Zhanylsyn U. Urasheva, Alima A. Khamidulla, Zhanylsyn N. Gaisiyeva, Gulnar B. Kabdrakhmanova, Aigul P. Yermagambetova, Aigerim B. Utegenova, Anastasiya G. Ishutina, Moldir M. Zhanuzakova, Moldir K. Omash. ANALYSIS OF RISK FACTORS FOR ISCHEMIC STROKE IN RURAL RESIDENTS OF THE AKTOBE REGION.	144-150
Bikbaeva Karina R, Kovalenko Elizaveta V, Vedeleva Ksenia V, Pichkurova Galina S, Maranyan Marina A, Baybuz Bogdan V, Baymurzaev Ibragim A, Cenko Evgeniy A, Kurmagomadov Adam A, Ataev Ahmed B, Malsagov Shahbulat Kh.-B. EVALUATION OF THE EFFECT OF REBAMIPIDE ON THE PROGRESSION OF ULCERATIVE COLITIS IN RATS IN THE EXPERIMENT.	151-153
Oleg Batiuk, Iryna Hora, Valeriy Kolesnyk, Inna Popovich, Oleksandr Sofilkanych. MEDICAL AND LEGAL ISSUES OF OBSERVING THE RIGHTS OF A PERSON WITH A MENTAL ILLNESS WHO HAS BECOME A PARTICIPANT IN CRIMINAL PROCEEDINGS.	154-160

SAFE USE OF INTRA-OPERATIVE TOURNIQUETS IN A DISTRICT HOSPITAL IN THE UK-AN AUDIT STUDY IN ORTHOPAEDIC THEATRES AND REVIEW OF CURRENT LITERATURE

Arnab Sain, Ralph Keita, Arunava Ray, Nauman Manzoor, Arsany Metry, Ahmed Elkilany, Kanishka Wattage, Michele Halasa, Jack Song Chia, Fahad Hussain, Odiamehi Aisabokhale, Zain Sohail, Vivek Deshmukh, Adhish Avasthi.

Department of Orthopaedics, Worthing Hospital, University Hospitals Sussex NHS Foundation Trust, United Kingdom

Abstract.

The use of tourniquet is common in orthopaedic surgeries as it reduces blood loss, enhances visualization of the operating field, and leads to quicker procedures. However, the use of tourniquet has certain risks which can be avoided by following guidelines like British Orthopaedic Association Standards for Trauma (BOAST) guidelines for safe use of tourniquet. This audit study was done in a District general hospital to check the compliance of two trauma theatres with BOAST guidelines. The audit found that there was poor documentation of tourniquet details in the operation notes (10%). Regarding tourniquet time and pressure, the compliance in the two theatres was 95 % & 97.5 %. The recommendations of this audit were to use a template to improve documentation of tourniquet details in the operation notes and training of theatre staff on BOAST guidelines for safe use of tourniquet.

Key words. Tourniquet, safe use guidelines, BOAST, orthopaedic trauma, theatres.

Introduction.

The tourniquet is an externally applied device to a body part to occlude the underlying blood flow and used in orthopaedic surgeries. The main advantage is reduced blood loss during surgery and also enhances the visualization of the surgical field and reduces operative time. For effective use of tourniquet, the tourniquet pressure should be greater than the arterial blood pressure of the limb [1,2]. The pressure required to stop arterial blood flow in a limb is called the limb occlusion pressure (LOP) or the arterial occlusion pressure (AOP). LOP is measured by serial inflation of pressure cuff until loss of arterial pulse of the limb on doppler ultrasound [2-4].

However, the use of tourniquet is associated with several risks such as nerve injuries (most common complication) [5], compartment syndrome [6], rhabdomyolysis [7], pressure sores, digital necrosis, chemical burns [8], deep vein thrombosis leading to pulmonary embolism [9], tourniquet pain [10].

The British Orthopaedic Association Standards for Trauma (BOAST) has set guidelines for safe use of intra-operative tourniquets, and it provides specific recommendations regarding safe limits of tourniquet pressure and duration. The BOAST guidelines also recommend clear documentation regarding skin site.

In our audit study, we aimed to check the compliance with BOAST guidelines during use of tourniquet in orthopaedic theatres in a District General Hospital in the UK [11].

Materials and Methods.

The study was a retrospective audit conducted in the orthopaedic theatres at Worthing Hospital. The audit was registered with the

hospital audit department with registration number 1661. The audit was a retrospective analysis of 40 patients each from two trauma theatres (Theatre 8 & Theatre9) from March 2022 till June 2022. The data was collected from theatre record books along with operation notes.

The data collected included patient name, age, sex, type of procedure and indication, the tourniquet pressure used and the tourniquet time and documentation of skin site. The study included only orthopaedic trauma patients and elective cases were excluded from the study.

Results.

In theatre 8, 21 patients had upper limb surgery with an average tourniquet time 45 minutes and 19 patients had lower limb surgery with an average tourniquet time of 56 minutes. In Theatre 9, 19 patients had upper limb surgery with an average tourniquet time of 54 minutes and 21 patients had lower limb surgery with an average tourniquet time of 73 minutes.

In theatre 8, 39 out of 40 patients had a tourniquet time less than 120 minutes (compliance rate of 97.5 percent). In one case where tourniquet time was exceeded there was no documentation of the tourniquet time in the operation note and no justification regarding the need to go above 120 minutes as per BOA guidelines.

In theatre 9, 38 out of 40 patients had a tourniquet time less than 120 minutes (compliance rate of 95 percent). There was a mention of tourniquet time being of 139 minutes for one of the operations, but no justification was given. The other operation where the tourniquet time was noted as 132 minutes in the theatre record book had some discrepancy with the operation note, where the tourniquet time was recorded as 95 minutes.

The results showed that there was poor documentation regarding the tourniquet time and duration in the operation notes as only 10 % of the operation notes had this information. The theatre record book contained most of the data.

With regards to compliance with BOAST guidelines with respect to tourniquet pressure and time, theatre 8 had 97.5 % compliance and theatre 9 had 95% compliance (Figure 1). In none of the theatres there was any documentation justifying the cause of increased duration of tourniquet time. Also, there was no documentation of tourniquet site.

Discussion and Review of Literature.

The use of tourniquet has led to less blood loss during limb surgery and also better visualization of the operative field leading to quicker procedures which is beneficial for the patient and improves outcomes. However, the use of tourniquet is associated with various complications like nerve injuries (most common complication) [5], compartment syndrome [6],




Figure 1. Compliance with BOAST guidelines with respect to tourniquet time in Theatre 8 and Theatre 9.

rhabdomyolysis [7], pressure sores, digital necrosis, chemical burns [8], deep vein thrombosis leading to pulmonary embolism [9], tourniquet pain [10].

The limb should be exsanguinated by simple elevation or by use of an Esmarch bandage or tourniquet exsanguinators which empties the blood vessels distally to the proximal part of the limb before inflation of the tourniquet. This creates a clear operating field, reduces blood loss, and decreases the risk of micro emboli at the time of release of tourniquet. Exsanguination provides a mode of autotransfusion of blood into the central circulation [1]. The optimum time and angle of elevation for maximum effect of exsanguinations of upper limb is 5 mins at 90 degrees [12] and for lower limb is 5 mins at 45 degrees.

After inflation of tourniquet there is progressive cellular hypoxia, acidosis and decrease in temperature of the occluded limb. The muscle is more susceptible to ischaemia than the nerve. Under tourniquet muscle has reduced pH and pO₂, increase in pCO₂, K⁺ and Lactate. There is evidence of tissue odema if tourniquet is used more than 60 mins [1]. Tourniquet pain develops in 66% patients after 30-60 mins of tourniquet use [13].

Before application of tourniquet, the pressure source, cuff, regulator, and tubing with connectors should be checked. Wider cuffs have better transmission of pressure during tissue compression and have lower chance of pressure related complications. Also, wider cuffs were less painful than narrower cuffs [14].

Documentation of tourniquet time and pressure is vital particularly in operation notes. The compliance with operation notes was variable in different studies. In two studies done in Hongkong and in India the documentation of tourniquet time and pressure in operation notes was poor with compliance of 32% [15] and 42 % [16]. Another study found that the introduction of an operation note template improved documentation of tourniquet time by 49 %. In our study the documentation of tourniquet details was poor with only 10%. As a recommendation on improvement in documentation of tourniquet details, an operation template is the best way to improve documentation.

Regarding compliance with BOAST guidelines with respect to tourniquet pressure and time, in study by Vayalapa et al showed compliance of 97 % in both their audit cycles [2]. In our study the compliance was 95% in theatre 9 and 97.5% in theatre 8. In cases where tourniquet time was exceeded no clear documentation was done either in operation notes or in theatre record book. In previous studies the main reason for this is lack of awareness regarding BOAST guidelines in theatre personnel. Also, there is a tendency to use standard tourniquet inflation pressure instead of calculating the pressure as per the systolic pressure. Studies have shown that there is a lack of formal training of staff on BOAST guidelines on safe use of tourniquet [2,17,18,19].

Conclusion.

The audit study concluded that there was lack of documentation on tourniquet details particularly in the operation notes. This problem can be effectively mitigated by using an operation template. In cases where the tourniquet time was exceeded there should be documentation justifying the need for that. The compliance with regards to tourniquet pressure and time was good and comparable with other studies, however that can be improved by proper training of staff on safe use of tourniquet. The lack of awareness can be solved by organizing regular training courses for theatre staff. Our audit study was a single cycle of an audit , so a reaudit after implementation of recommendations is necessary.

REFERENCES

1. Sharma J.P, Salhotra R. Tourniquets in orthopedic surgery. Indian journal of orthopaedics. 2012;46:377-383.
2. Vayalapa S, Guerero D.N, Balasubramanian B, et al. Improving the Safety of Tourniquet Use in a Trauma Theatre According to the British Orthopaedic Association Guidelines: A Closed Loop Audit. Cureus. 2024;16:e51601.
3. Kasem SA, Bassiouny AA, Rashwan DA, et al. Minimal inflation tourniquet pressure using induced hypotension with limb occlusion pressure determination or arterial occlusion pressure estimation in upper limb surgery: a randomized double-blinded comparative study. Anesth Pain Med. 2020;10:e102124.

4. Deloughry JL, Griffiths R. Arterial tourniquets. Continuing education in anaesthesia, critical care and pain. 2009;9:56-60.
5. Odinson A, Finsen V. Tourniquet use and its complications in Norway. *J Bone Joint Surg Br.* 2006;88:1090-2.
6. Hirvensalo E, Tuominen H, Lapinsuo M, et al. Compartment syndrome of the lower limb caused by a tourniquet: a report of two cases. *J Orthop Trauma.* 1992;6:469-72.
7. Türkmen İ, Esenkaya İ, Unay K, et al. Rhabdomyolysis after tourniquet use in proximal tibial osteotomy: a case report and review of the literature. *Acta Orthop Traumatol Turc.* 2015;49:338-41.
8. Yang JH, Lim H, Yoon JR, et al. Tourniquet associated chemical burn. *Indian J Orthop.* 2012;46:356-9.
9. Song JE, Chun DH, Shin JH, et al. Pulmonary thromboembolism after tourniquet inflation under spinal anesthesia—a case report. *Korean J Anesthesiol.* 2010;59:S82-5.
10. Kamath K, Kamath SU, Tejaswi P. Incidence and factors influencing tourniquet pain. *Chin J Traumatol.* 2021;24:291-4.
11. BOAST - the safe use of intraoperative tourniquets. 2021.
12. Warren PJ, Hardiman PJJ-P, Woolf VJ. Limb exsanguination. I. The arm: effect of angle of elevation and arterial compression. *Ann R Coll Surg Engl.* 1992;74:320-2.
13. Micheal K. Urban. Anaesthesia for orthopaedic surgery. In: Miller RD, editor. *Miller's Anesthesia.* 7th ed. Philadelphia: Churchill Livingstone Publishers; 2010:2252.
14. Estebe J.P, Le Naoures A, Chemaly L, et al. Tourniquet pain in a volunteer study: effect of changes in cuff width and pressure. *Anaesthesia.* 2000;55:21-26.
15. Sweed TA, Bonajmah AA, Mussa MA. Audit of operation notes in an orthopaedic unit. *J Orthop Surg (Hong Kong).* 2014;22:218-20.
16. Tuteja S, Tiwari A, Bhanushali J, et al. Results of an audit of orthopaedic operation notes from a tertiary care centre: are we doing it right and can we do more?. *Indian J Orthop.* 2022;56:2223-7.
17. Younger AS, Kalla TP, McEwen JA, et al. Survey of tourniquet use in orthopaedic foot and ankle surgery. *Foot Ankle Int.* 2005;26:208-17.
18. Kalla TP, Younger A, McEwen JA, et al. Survey of tourniquet use in podiatric surgery. *J Foot Ankle Surg.* 2003;42:68-76.
19. Sadri A, Braithwaite IJ, Abdul-Jabar HB, et al. Understanding of intra-operative tourniquets amongst orthopaedic surgeons and theatre staff—a questionnaire study. *Ann R Coll Surg Engl.* 2010;92:243-5.