GEORGIAN MEDICAL MEWS

ISSN 1512-0112

NO 7-8 (352-353) Июль-Август 2024

ТБИЛИСИ - NEW YORK



ЕЖЕМЕСЯЧНЫЙ НАУЧНЫЙ ЖУРНАЛ

Медицинские новости Грузии საქართველოს სამედიცინო სიახლენი

GEORGIAN MEDICAL NEWS

Monthly Georgia-US joint scientific journal published both in electronic and paper formats of the Agency of Medical Information of the Georgian Association of Business Press. Published since 1994. Distributed in NIS, EU and USA.

GMN: Georgian Medical News is peer-reviewed, published monthly journal committed to promoting the science and art of medicine and the betterment of public health, published by the GMN Editorial Board since 1994. GMN carries original scientific articles on medicine, biology and pharmacy, which are of experimental, theoretical and practical character; publishes original research, reviews, commentaries, editorials, essays, medical news, and correspondence in English and Russian.

GMN is indexed in MEDLINE, SCOPUS, PubMed and VINITI Russian Academy of Sciences. The full text content is available through EBSCO databases.

GMN: Медицинские новости Грузии - ежемесячный рецензируемый научный журнал, издаётся Редакционной коллегией с 1994 года на русском и английском языках в целях поддержки медицинской науки и улучшения здравоохранения. В журнале публикуются оригинальные научные статьи в области медицины, биологии и фармации, статьи обзорного характера, научные сообщения, новости медицины и здравоохранения. Журнал индексируется в MEDLINE, отражён в базе данных SCOPUS, PubMed и ВИНИТИ РАН. Полнотекстовые статьи журнала доступны через БД EBSCO.

GMN: Georgian Medical News – საქართველოს სამედიცინო სიახლენი – არის ყოველთვიური სამეცნიერო სამედიცინო რეცენზირებადი ჟურნალი, გამოიცემა 1994 წლიდან, წარმოადგენს სარედაქციო კოლეგიისა და აშშ-ის მეცნიერების, განათლების, ინდუსტრიის, ხელოვნებისა და ბუნებისმეტყველების საერთაშორისო აკადემიის ერთობლივ გამოცემას. GMN-ში რუსულ და ინგლისურ ენებზე ქვეყნდება ექსპერიმენტული, თეორიული და პრაქტიკული ხასიათის ორიგინალური სამეცნიერო სტატიები მედიცინის, ბიოლოგიისა და ფარმაციის სფეროში, მიმოხილვითი ხასიათის სტატიები.

ჟურნალი ინდექსირებულია MEDLINE-ის საერთაშორისო სისტემაში, ასახულია SCOPUS-ის, PubMed-ის და ВИНИТИ РАН-ის მონაცემთა ბაზებში. სტატიების სრული ტექსტი ხელმისაწვდომია EBSCO-ს მონაცემთა ბაზებიდან.

WEBSITE

www.geomednews.com

К СВЕДЕНИЮ АВТОРОВ!

При направлении статьи в редакцию необходимо соблюдать следующие правила:

- 1. Статья должна быть представлена в двух экземплярах, на русском или английском языках, напечатанная через полтора интервала на одной стороне стандартного листа с шириной левого поля в три сантиметра. Используемый компьютерный шрифт для текста на русском и английском языках Times New Roman (Кириллица), для текста на грузинском языке следует использовать AcadNusx. Размер шрифта 12. К рукописи, напечатанной на компьютере, должен быть приложен CD со статьей.
- 2. Размер статьи должен быть не менее десяти и не более двадцати страниц машинописи, включая указатель литературы и резюме на английском, русском и грузинском языках.
- 3. В статье должны быть освещены актуальность данного материала, методы и результаты исследования и их обсуждение.

При представлении в печать научных экспериментальных работ авторы должны указывать вид и количество экспериментальных животных, применявшиеся методы обезболивания и усыпления (в ходе острых опытов).

- 4. К статье должны быть приложены краткое (на полстраницы) резюме на английском, русском и грузинском языках (включающее следующие разделы: цель исследования, материал и методы, результаты и заключение) и список ключевых слов (key words).
- 5. Таблицы необходимо представлять в печатной форме. Фотокопии не принимаются. Все цифровые, итоговые и процентные данные в таблицах должны соответствовать таковым в тексте статьи. Таблицы и графики должны быть озаглавлены.
- 6. Фотографии должны быть контрастными, фотокопии с рентгенограмм в позитивном изображении. Рисунки, чертежи и диаграммы следует озаглавить, пронумеровать и вставить в соответствующее место текста в tiff формате.

В подписях к микрофотографиям следует указывать степень увеличения через окуляр или объектив и метод окраски или импрегнации срезов.

- 7. Фамилии отечественных авторов приводятся в оригинальной транскрипции.
- 8. При оформлении и направлении статей в журнал МНГ просим авторов соблюдать правила, изложенные в «Единых требованиях к рукописям, представляемым в биомедицинские журналы», принятых Международным комитетом редакторов медицинских журналов http://www.spinesurgery.ru/files/publish.pdf и http://www.nlm.nih.gov/bsd/uniform_requirements.html В конце каждой оригинальной статьи приводится библиографический список. В список литературы включаются все материалы, на которые имеются ссылки в тексте. Список составляется в алфавитном порядке и нумеруется. Литературный источник приводится на языке оригинала. В списке литературы сначала приводятся работы, написанные знаками грузинского алфавита, затем кириллицей и латиницей. Ссылки на цитируемые работы в тексте статьи даются в квадратных скобках в виде номера, соответствующего номеру данной работы в списке литературы. Большинство цитированных источников должны быть за последние 5-7 лет.
- 9. Для получения права на публикацию статья должна иметь от руководителя работы или учреждения визу и сопроводительное отношение, написанные или напечатанные на бланке и заверенные подписью и печатью.
- 10. В конце статьи должны быть подписи всех авторов, полностью приведены их фамилии, имена и отчества, указаны служебный и домашний номера телефонов и адреса или иные координаты. Количество авторов (соавторов) не должно превышать пяти человек.
- 11. Редакция оставляет за собой право сокращать и исправлять статьи. Корректура авторам не высылается, вся работа и сверка проводится по авторскому оригиналу.
- 12. Недопустимо направление в редакцию работ, представленных к печати в иных издательствах или опубликованных в других изданиях.

При нарушении указанных правил статьи не рассматриваются.

REQUIREMENTS

Please note, materials submitted to the Editorial Office Staff are supposed to meet the following requirements:

- 1. Articles must be provided with a double copy, in English or Russian languages and typed or computer-printed on a single side of standard typing paper, with the left margin of 3 centimeters width, and 1.5 spacing between the lines, typeface Times New Roman (Cyrillic), print size 12 (referring to Georgian and Russian materials). With computer-printed texts please enclose a CD carrying the same file titled with Latin symbols.
- 2. Size of the article, including index and resume in English, Russian and Georgian languages must be at least 10 pages and not exceed the limit of 20 pages of typed or computer-printed text.
- 3. Submitted material must include a coverage of a topical subject, research methods, results, and review.

Authors of the scientific-research works must indicate the number of experimental biological species drawn in, list the employed methods of anesthetization and soporific means used during acute tests.

- 4. Articles must have a short (half page) abstract in English, Russian and Georgian (including the following sections: aim of study, material and methods, results and conclusions) and a list of key words.
- 5. Tables must be presented in an original typed or computer-printed form, instead of a photocopied version. Numbers, totals, percentile data on the tables must coincide with those in the texts of the articles. Tables and graphs must be headed.
- 6. Photographs are required to be contrasted and must be submitted with doubles. Please number each photograph with a pencil on its back, indicate author's name, title of the article (short version), and mark out its top and bottom parts. Drawings must be accurate, drafts and diagrams drawn in Indian ink (or black ink). Photocopies of the X-ray photographs must be presented in a positive image in **tiff format**.

Accurately numbered subtitles for each illustration must be listed on a separate sheet of paper. In the subtitles for the microphotographs please indicate the ocular and objective lens magnification power, method of coloring or impregnation of the microscopic sections (preparations).

- 7. Please indicate last names, first and middle initials of the native authors, present names and initials of the foreign authors in the transcription of the original language, enclose in parenthesis corresponding number under which the author is listed in the reference materials.
- 8. Please follow guidance offered to authors by The International Committee of Medical Journal Editors guidance in its Uniform Requirements for Manuscripts Submitted to Biomedical Journals publication available online at: http://www.nlm.nih.gov/bsd/uniform_requirements.html http://www.icmje.org/urm_full.pdf
- In GMN style for each work cited in the text, a bibliographic reference is given, and this is located at the end of the article under the title "References". All references cited in the text must be listed. The list of references should be arranged alphabetically and then numbered. References are numbered in the text [numbers in square brackets] and in the reference list and numbers are repeated throughout the text as needed. The bibliographic description is given in the language of publication (citations in Georgian script are followed by Cyrillic and Latin).
- 9. To obtain the rights of publication articles must be accompanied by a visa from the project instructor or the establishment, where the work has been performed, and a reference letter, both written or typed on a special signed form, certified by a stamp or a seal.
- 10. Articles must be signed by all of the authors at the end, and they must be provided with a list of full names, office and home phone numbers and addresses or other non-office locations where the authors could be reached. The number of the authors (co-authors) must not exceed the limit of 5 people.
- 11. Editorial Staff reserves the rights to cut down in size and correct the articles. Proof-sheets are not sent out to the authors. The entire editorial and collation work is performed according to the author's original text.
- 12. Sending in the works that have already been assigned to the press by other Editorial Staffs or have been printed by other publishers is not permissible.

Articles that Fail to Meet the Aforementioned Requirements are not Assigned to be Reviewed.

ᲐᲕᲢᲝᲠᲗᲐ ᲡᲐᲧᲣᲠᲐᲓᲦᲔᲑᲝᲓ!

რედაქციაში სტატიის წარმოდგენისას საჭიროა დავიცვათ შემდეგი წესები:

- 1. სტატია უნდა წარმოადგინოთ 2 ცალად, რუსულ ან ინგლისურ ენებზე,დაბეჭდილი სტანდარტული ფურცლის 1 გვერდზე, 3 სმ სიგანის მარცხენა ველისა და სტრიქონებს შორის 1,5 ინტერვალის დაცვით. გამოყენებული კომპიუტერული შრიფტი რუსულ და ინგლისურენოვან ტექსტებში Times New Roman (Кириллица), ხოლო ქართულენოვან ტექსტში საჭიროა გამოვიყენოთ AcadNusx. შრიფტის ზომა 12. სტატიას თან უნდა ახლდეს CD სტატიით.
- 2. სტატიის მოცულობა არ უნდა შეადგენდეს 10 გვერდზე ნაკლებს და 20 გვერდზე მეტს ლიტერატურის სიის და რეზიუმეების (ინგლისურ,რუსულ და ქართულ ენებზე) ჩათვლით.
- 3. სტატიაში საჭიროა გაშუქდეს: საკითხის აქტუალობა; კვლევის მიზანი; საკვლევი მასალა და გამოყენებული მეთოდები; მიღებული შედეგები და მათი განსჯა. ექსპერიმენტული ხასიათის სტატიების წარმოდგენისას ავტორებმა უნდა მიუთითონ საექსპერიმენტო ცხოველების სახეობა და რაოდენობა; გაუტკივარებისა და დაძინების მეთოდები (მწვავე ცდების პირობებში).
- 4. სტატიას თან უნდა ახლდეს რეზიუმე ინგლისურ, რუსულ და ქართულ ენებზე არანაკლებ ნახევარი გვერდის მოცულობისა (სათაურის, ავტორების, დაწესებულების მითითებით და უნდა შეიცავდეს შემდეგ განყოფილებებს: მიზანი, მასალა და მეთოდები, შედეგები და დასკვნები; ტექსტუალური ნაწილი არ უნდა იყოს 15 სტრიქონზე ნაკლები) და საკვანძო სიტყვების ჩამონათვალი (key words).
- 5. ცხრილები საჭიროა წარმოადგინოთ ნაბეჭდი სახით. ყველა ციფრული, შემაჯამებელი და პროცენტული მონაცემები უნდა შეესაბამებოდეს ტექსტში მოყვანილს.
- 6. ფოტოსურათები უნდა იყოს კონტრასტული; სურათები, ნახაზები, დიაგრამები დასათაურებული, დანომრილი და სათანადო ადგილას ჩასმული. რენტგენოგრამების ფოტოასლები წარმოადგინეთ პოზიტიური გამოსახულებით tiff ფორმატში. მიკროფოტო-სურათების წარწერებში საჭიროა მიუთითოთ ოკულარის ან ობიექტივის საშუალებით გადიდების ხარისხი, ანათალების შეღებვის ან იმპრეგნაციის მეთოდი და აღნიშნოთ სუ-რათის ზედა და ქვედა ნაწილები.
- 7. სამამულო ავტორების გვარები სტატიაში აღინიშნება ინიციალების თანდართვით, უცხოურისა უცხოური ტრანსკრიპციით.
- 8. სტატიას თან უნდა ახლდეს ავტორის მიერ გამოყენებული სამამულო და უცხოური შრომების ბიბლიოგრაფიული სია (ბოლო 5-8 წლის სიღრმით). ანბანური წყობით წარმოდგენილ ბიბლიოგრაფიულ სიაში მიუთითეთ ჯერ სამამულო, შემდეგ უცხოელი ავტორები (გვარი, ინიციალები, სტატიის სათაური, ჟურნალის დასახელება, გამოცემის ადგილი, წელი, ჟურნალის №, პირველი და ბოლო გვერდები). მონოგრაფიის შემთხვევაში მიუთითეთ გამოცემის წელი, ადგილი და გვერდების საერთო რაოდენობა. ტექსტში კვადრატულ ფჩხილებში უნდა მიუთითოთ ავტორის შესაბამისი N ლიტერატურის სიის მიხედვით. მიზანშეწონილია, რომ ციტირებული წყაროების უმეტესი ნაწილი იყოს 5-6 წლის სიღრმის.
- 9. სტატიას თან უნდა ახლდეს: ა) დაწესებულების ან სამეცნიერო ხელმძღვანელის წარდგინება, დამოწმებული ხელმოწერითა და ბეჭდით; ბ) დარგის სპეციალისტის დამოწმებული რეცენზია, რომელშიც მითითებული იქნება საკითხის აქტუალობა, მასალის საკმაობა, მეთოდის სანდოობა, შედეგების სამეცნიერო-პრაქტიკული მნიშვნელობა.
- 10. სტატიის ბოლოს საჭიროა ყველა ავტორის ხელმოწერა, რომელთა რაოდენობა არ უნდა აღემატებოდეს 5-ს.
- 11. რედაქცია იტოვებს უფლებას შეასწოროს სტატია. ტექსტზე მუშაობა და შეჯერება ხდება საავტორო ორიგინალის მიხედვით.
- 12. დაუშვებელია რედაქციაში ისეთი სტატიის წარდგენა, რომელიც დასაბეჭდად წარდგენილი იყო სხვა რედაქციაში ან გამოქვეყნებული იყო სხვა გამოცემებში.

აღნიშნული წესების დარღვევის შემთხვევაში სტატიები არ განიხილება.

Содержание:

Yevchuk YuI, Rozhko MM, Pantus AV, Yarmoshuk IR, Pantus PV. ANALYSIS OF THE CLINICAL EFFECTIVENESS OF USING THE CREATED COMBINED FIBRIN-BONE SCAFFOLD FOR THE RECONSTRUCTION OF BONE TISSUE DEFECTS OF THE JAWS
Anton Yu. Postnov, Tatiana V. Kirichenko, Yuliya V. Markina, Petr V. Chumachenko, Andrey V. Suslov, Alexandra G. Ivanova, Eduard R. Charchyan, Alexander M. Markin. INFLAMMATORY FACTORS IN DISSECTION OF THORACIC AORTIC ANEURYSM
Gohar Arajyan, Qristine Navoyan, Nvard Pahutyan, Hovhannes Hunanyan, Anahit Pogosyan, Hrachik Gasparyan. COMPREHENSIVE STUDY OF ANTIOXIDANT ACTIVITY OF OXALIC ACID DIAMIDE DERIVATIVES AND THEIR EFFECT ON THE CONCENTRATION OF MALONIC DIALDEHYDE IN THE BRAIN AND LIVER TISSUES OF WHITE RATS
Nino Abesadze, Jenaro Kristesashvili, Arsen Gvenetadze. LOW 25OHD IN ENDOMETRIOSIS- RISK FACTOR OR CONSEQUENCE?!
Stepanyan L, Lalayan G. STRESS RESILIENCE AND DECISION-MAKING UNDER PRESSURE: ENHANCING ATHLETIC PERFORMANCE IN COMPETITIVE SPORTS
Hasan M. Abed, Abdulameer M. Hussein, Sabah N. Jaber. ENDOVASCULAR INTERVENTIONS: A NEW INSIGHTS AND CLINICAL PRACTICE
Changsheng He, Jian Liu, Linhai Xu, Fanhua Sun, Yan Wang, Jia Lou. THE RELATIONSHIP BETWEEN SERUM INFLAMMATORY CYTOKINES AND HYPERLIPIDEMIC ACUTE PANCREATITIS47-49
Artemov O.V, Lytvynenko M.V, Chumachenko I.V, Bondarenko A.V, Dotsenko N.V, Ostapchuk K.V, Koshelnyk O.L, Gargin V.V. THE INFLUENCE OF THE DEMODEX MITE ON THE MORPHOLOGICAL PICTURE OF EYELID PAPILLOMA50-54
Othman K.M. Al-Sawaf, Mahmoud AM Fakhri. CHARACTERIZATION OF SERUM SERINE PROTEASE BIOCHEMICAL PROFILE IN PATIENTS WITH RENAL FAILURE55-58
Sergey Lee, Marat Assimov, Yuriy Ignatiev, Fatima Bagiyarova, Gulbanu Absatarova, Aizhan Kudaibergenova, Sholpan Mardanova, Tatyana Tsapenko, Baimakhan Tanabayev, Assel Ibrayeva, Anel Ibrayeva, Ildar Fakhradiyev. PREVALENCE AND FACTORS OF PROFESSIONAL BURNOUT AMONG PRIMARY HEALTHCARE WORKERS IN THE REPUBLIC OF KAZAKHSTAN: RESULTS OF A NATIONAL STUDY
I.A. Yusubov. RESULTS OF PERCUTANEOUS TREATMENT OF LIMITED FLUID FORMATIONS AFTER ABDOMINAL SURGERY
Nawar M. Abd-alaziz, Ammar L. Hussein, Mohammed M Abdul-Aziz. STUDY THE RELATIONSHIP BETWEEN OSTEOPROTEGERIN AND KIDNEY INJURY MOLECULE-1 AND SOME BIOCHEMICAL VARIABLES IN PATIENTS WITH KIDNEY STONES
Tsisana Giorgadze, Tinatin Gognadze. SUBSTRATE SPECIFICITY OF β-GLUCOSIDASE FROM YUCCA GLORIOSA LEAVES
Sheishenov Zhalil, Kemelbekov Kanatzhan, Joshibaev Seitkhan, Turtabaev Baglan, Zhunissov Bakhytzhan. COMPARATIVE ANALYSIS OF THE CLINICAL RESULTS OF PATIENTS WITH ASD OPERATED VIA RIGHT ANTERIOR MINITHORACOTOMY AND MEDIAN STERNOTOMY
Sosonna L, Ohurtsov O, Piriatinska N, Vdovitchenko V, Seleznova R, Kolba O, Gryzodub D, Rozhkova O, Shevtsov O. INDIVIDUAL ANATOMICAL VARIABILITY OF THE SKULL'S FACIAL SECTION CONSIDERING GENDER AND CRANIOTYPE BASED ON COMPUTED TOMOGRAPHY DATA
Osminina M.K, Aslamazova A.E, Podchernyaeva N.S, Khachatryan L.G, Velikoretskaya M.D, Chebysheva S.N, Polyanskaya A.V. SYSTEMIC OR LIMITED IS HEMISCLERODERMA OF FACE IN A PERSON WITH UVEITIS? EXPERIENCE OF 10 CASES OF UVEITIS IN HEMISCLERODERMA OF FACE FROM ONE RHEUMATOLOGY CENTER
F.T. Khalilova, A.A. Kerimov. CLINICAL AND LABORATORY CHARACTERISTICS OF THE LATENT FORM OF POLYCYTHEMIA VERA101-105
Ahlam S. Ibrahim, Sukayna H. Rashed. ISOLATION AND PURIFICATION OF TRANSGLUTAMINASE 1 USING BIOCHEMICAL TECHNIQUES
Tingting Li, Xu Zhang, Baohong Xue, Lianping He, Qiaoqiao Chen, Dexun Zhao. THE RELATIONSHIP BETWEEN MENTAL HEALTH AND PHYSICAL ACTIVITY AMONG STUDENTS FROM A PRIVATE UNIVERSITY: A CROSS-SECTION STUDY
Narkhojayev Nurgali, Turmetov Ibadulla, Kemelbekov Kanatzhan, Bektayev Erkebai, Akhmetov Almasbek, Zhunissov Bakhytzhan. RESULTS OF SURGICAL TREATMENT OF PECTUS EXCAVATUM IN CHILDREN AND ADOLESCENTS

Krushelnytska HL, Batryn OV, Ryzhenko LM, Lytvyn NA, Dobrianska NV, Lyga AI. INFORMATION FACTORS OF MEDIA INFLUENCE ON THE FORMATION OF STATE POLICY IN THE FIELD OF LEGAL REGULATION OF BIOMEDICAL TECHNOLOGIES
Vahe Ashot Ter-Minasyan. EVALUATION OF KNOWLEDGE AND ATTITUDE REGARDING CERVICAL CANCER SCREENING PRACTICE: A MULTICENTER REGIONAL STUDY
Muhsin S.G. Almozie'l, Abbas A. Khudhair, Falah Hassan Shari. REMEDIAL INTERVENTION OF FERTILITY AGENT AND GENE 35 ON INDUCED CYSTIC OVARY IN RATS
Rongzheng Yuan, Hui Wang, Jing Chen. THE EFFECT OF LOW MOLECULAR WEIGHT HEPARIN SODIUM IN THE TREATMENT OF ACUTE EXACERBATION OF CHRONIC OBSTRUCTIVE PULMONARY DISEASE COMORBID WITH PULMONARY HEART DISEASE ON PROMOTING THE BALANCE OF BLOOD VESSELS
Arailym Maikenova, Alexander Nersesov, Elmira Kuantay, Mukhtar Kulimbet, Massimo Giuseppe Colombo, Chavdar Pavlov, Yerkezhan
Yerlanova. EVALUATION OF PREDICTORS OF INEFFECTIVENESS OF ANTIVIRAL THERAPY FOR CHRONIC HEPATITIS C IN THE REPUBLIC OF KAZAKHSTAN: A MATCHED CASE-CONTROL STUDY
Ahmed N. Ali, Muna A. Kashmoola. EVALUATION OF PROTEIN C AND S IN β-THALASSEMIA MAJOR
Sh.Tsiklauri, N.Nakudashvili, M.Lomaia. EFFECT OF INTRANASAL ELECTROPHORESIS WITH 5% POTASSIUM IODATE SOLUTION ON CLINICAL OUTCOME OF PATIENTS WITH HYPERTROPHIC RHINITIS
Fang Xu, Zhijuan Xu, Ming Li. INTRAVITREAL INJECTION CONBERCEPT IMPROVES THE BEST-CORRECTED VISUAL ACUITY IN PATIENTS WITH WET AGE- RELATED MACULAR EDEMA
Lilit Darbinyan, Margarita Danielyan, Vergine Chavushyan, Karen Simonyan, Michael Babakhanyan, Lilia Hambardzumyan, Larisa Manukyan, Kristine Karapetyan, Lusya Hovhannisyan. THE PROTECTIVE EFFECTS OF SELENIUM-ENRICHED HYDROPONIC RADISH ON PARACETAMOL-INDUCED LIVER DAMAGE IN RATS
Grygorova A.O, Grygorov S.M, Yaroslavska Yu.Yu, Mykhailenko N.M, Demyanyk D.S, Steblianko A.O, Rak O.V, Voloshan O.O, Nazaryan R.S.
SIGNS OF ORAL CAVITY MICROCIRCULATORY DISORDERS IN ADOLESCENTS WHO SMOKE
Ali H. Kadhim, Nihad N. Hilal, Taghreed AH. Nassir. A COMPARATIVE STUDY ON THE VARIABLE EFFECTS OF ALCOHOL AND NON-ALCOHOL-RELATED FATTY LIVER DISEASE ON METABOLIC AND INFLAMMATORY BIOMARKERS
Papoyan Varduhi, Galstyan Alina, Sargsyan Diana. FACTOR ANALYSIS OF THE COMPETENCIES OF PERSONAL RESOURCES OF SPECIALIST
Chulpanov Utkir, Turdaliyeva Botagoz, Buleshov Myrzatai, Zhanabaev Nurlan, Kanatzhan Kemelbekov. COMPARATIVE EVALUATION OF THE EFFECTIVENESS OF INNOVATIVE HIGH-TECH CARDIAC SURGERY IN PATIENTS WHO HAVE SUFFERED AN ACUTE MYOCARDIAL INFARCTION
Tea Charkviani, Jenara Kristasashvili, Tamar Barbakadze, Mariam Gabadze, Tamar Kbilashvili, Mariam Makharadze. THE RELATIONSHIP BETWEEN FOLLICLE SIZE, OOCYTE MATURATION, BLASTOCYST FORMATION, BLASTOCYST PLOIDY, AND PREGNANCY OUTCOMES IN YOUNG WOMEN UNDERGOING IVF
Yunfei Wu, Koulong Wu, TianhuaDu. STUDY ON THE EFFECTS OF ART PAINTING COMBINED WITH SPORTS ON MYOPIA PREVENTION AND VISION IMPROVEMENT
Lulëjeta Ferizi-Shabani, Shefqet Mrasori, Valbona Ferizi, Gonxhe Barku, Milazim Gjocaj, Blerim Krasniqi, Basri Lenjani. EVALUATION OF DENTAL AND PERIODONTAL STATUS IN CHILDREN WITH TYPE 1 DIABETES MELLITUS208-212
Rana Dawood Salman Al-kamil, Mustafa Ragheb Abed, Sanaryh Mohammed Al-awad, H. N. K. AL-Salman, Hussein H. Hussein, Dawood Chaloob Hilyail, Falah Hassan Shari.
Chalood Filyan, Palan Flassan Shari. ISOLATION, CHARACTERIZATION, AND ANTIHYPERTENSIVE ACTIVITY ALKALOIDS EXTRACTED FROM THE LEAVES OF THE ALSTONIA SCHOLARIS PLANT
Tchernev G, Broshtilova V, Kordeva S. SHARK PEDICLE ISLAND FLAP FOR BASAL CELL CARCINOMA OF THE PERIALAR ZONE OF THE NOSE: PHOTOXICITY AND PHOTOCARCINOGENICITY MEDIATED BY POTENTIALLY NITROSAMINE CONTAMINATED DRUG INTAKE -A NEW EXPLANATION FOR THE SKIN CANCERS PATHOGENESIS?

Meruert T. Orazgalieva, Meyrbek J. Aimagambetov, Zhanna D. Bryzhakhina, Serik D. Zhanybekov, Ainash S. Orazalina. RISK FACTORS FOR THE DEVELOPMENT OF COAGULOPATHY DURING SURGERY IN MECHANICAL JAUNDICE223-228
Noor N. Noori, Nawal A. Murtadha. UNCONTROLLED TYPE 2 DIABETES MELLITUS MODULATED PLASMA LEVELS OF LIPID CATABOLIC PROTEINS229-233
Ling-Ling Zhou, Zhou-Zhou Lin, Lian-Ping He. PREVALENCE OF DEPRESSION AMONG UNIVERSITY STUDENTS IN CHINA: A PROTOCOL FOR A SYSTEMATIC REVIEW AND META-ANALYSIS
Nadine Khayyat, Sima Kalaldeh, Suha Khalifa. OPTIMISING THE CLINICAL ASSESSMENT OF CHILDHOOD AND ADOLESCENT OBESITY IN JORDAN
Shuasheva Y.A, Buleshov M.A, Kemelbekov K.S. CLINICAL, IMMUNOLOGICAL AND THESIOGRAPHIC CHARACTERISTICS RHEUMA-TOID ARTHRITIS AND CHRONIC RHEUMATICHEART DISEASE
Sana A. Abdulmawjood, Eman S. Mahmoud, Rana T Altaee. ASSESSMENT OF CIPROFLOXACIN EFFECTS ON SOME CHICKS' ORGANS: A COMPREHENSIVE BIOCHEMICAL AND HISTOLOGICALSTUDY
Knarik V. Kazaryan, Naira G. Hunanyan, Margarita H. Danielyan, Rosa G. Chibukchyan, Yulia Y. Trofimova, Arus V. Mkrtchyan, Kristine V. Karapetyan, Karwan H. Syan, Tatevik A. Piliposyan. REGULATION OF SPONTANEOUS ELECTRICAL ACTIVITY IN THE ORGANS OF RE-PRODUCTIVE SYSTEM BY OXYTOCIN
Lantukh I.V, Kucheriavchenko V.V, Yurko K.V, Bondarenko A.V, Merkulova N.F, Mohylenets O.I, Gradil G.I, Bondar O.Ye, Bodnia I.P, Burma Ya.I, Tsyko O.V, Tkachenko V.G. PSYCHOLOGICAL FEATURES OF REHABILITATION OF HIV-INFECTED PATIENTS
Serikbayeva Saltanat, Shaimerdenova Gulbanu, Ormanov Namazbai, Ormanov Talgat, Abuova Gulzhan, Kaishibayeva Gulnaz, Kemelbekov Kanatzhan. PEROXIDATION OF SALIVA LIPIDS IN PATIENTS WITH POSTCOVID SYNDROME DURING HIRUDOTHERAPY265-269
M.V. Poghosyan, H.Y. Stepanyan, Avetisyan Z.A, J.S. Sarkissian. THE EFFECTS OF HYDROCORTISONE ON SYNAPTIC PROCESSES IN PARKINSON'S DISEASE UNDERLYING THE POTENTIAL THERAPEUTICSTRATEGIES
Changsheng He, Jian Liu, Linhai Xu, Fanhua Sun. THE EFFECT OF PERCUTANEOUS CATHETER DRAINAGE COMBINED WITH SOMATOSTATIN ON INFLAMMATION AND PLASMA THROMBOXANE 2, PROSTACYCLIN 12 LEVELS IN PATIENTS WITH SEVERE PANCREATITIS
Tea Chitadze, Nino Sharashidze, Tamar Rukhadze, Nino Lomia, Giorgi Saatashvili. EVALUATION OF LEFT VENTRICULAR SYSTOLIC FUNCTION IN POSTMENOPAUSAL WOMEN WITH BREAST CANCER RECEIVING ADJUVANT ANTHRACYCLINE AND TRASTUZUMAB THERAPY: A 2-YEAR FOLLOW-UP STUDY284-293

PREVALENCE OF DEPRESSION AMONG UNIVERSITY STUDENTS IN CHINA: A PROTOCOL FOR A SYSTEMATIC REVIEW AND META-ANALYSIS

Ling-Ling Zhou¹, Zhou-Zhou Lin², Lian-Ping He^{1*}.

¹School of medicine, Taizhou university, Jiaojiang 318000, Zhejiang Province, China. ²School of clinical, Hainan medical university, Haikou 571199, Hainan Province, China.

Abstract.

Background: Depression is a mental disorder characterized by a persistent and long-lasting depressive mood.

Objective: The aim of this study was to analyze the prevalence of depression among university students in China and to provide guidance for psychological intervention and improvement of depression among this specific population. Methods: A systematic search was conducted on PubMed, CNKI, Wangfang database, and Web of Science to identify relevant articles. The articles were cross-sectional studies focusing on the prevalence of depression among university students in China, with clearly defined criteria for diagnosing depression included. Excel was used independently and in duplicate to perform the screening process that involved title, abstract screening, full-text screening, data extraction and assessment of risk of bias. The standards recommended by the Agency for Healthcare Research and Quality (AHRQ) was used to evaluate the quality of observational studies.

Results: STATA 11.0 was used to pool the outcomes and perform meta-analysis, including the prevalence of depression among university students and influential factors such as sex, major, region, and the impact of COVID-19.

Conclusion: Practical psychological interventions are needed in universities in China. Continuously attention is suggested to be paid to Chinese university students' mental health, especially after public health event such as COVID-19.

Key words. University students, depression, prevalence, meta-analysis, China.

Introduction.

Depression is a mental disorder characterized by a persistent and long-lasting depressive mood [1]. Depression is often accompanied by distinct and persistent depressive syndrome. Depression commonly manifests with physical symptoms such as fatigue, pain, or sleep disturbances, significantly affecting both the physical and mental well-being of individuals [2]. In contrast, a depressive mood is a normal change in mood that is temporary and does not have an impact on physical health or social relationships. However, depression can cause problems at school and work [3]. The performance of patients with depression can be divided into three aspects: Core symptom group, psychological symptom group, and physical symptom group. According to World Health Organization (WHO), an estimated 3.8% of the population experience depression, including 5% of adults (4% among men and 6% among women) and 5.7% of adults older than 60 years. This equates to around 280 million people worldwide who suffer from depression, highlighting its status as a significant public health issue [4].

Due to their immature physical and mental development, weak self-regulation, and self-control ability, university students face a multitude of self-and social problems. These challenges are further compounded by the heavy learning tasks they must undertake, which often result in strong psychological conflicts. Consequently, university students frequently experience depression and other negative moods. Depression in university students has been linked to various adverse outcomes, including lower academic performance, impaired roles, increased risk of dropping out, heightened anxiety levels, physical illness, reduced physical activity, unsafe sexual behavior, increased smoking, alcohol, and drug dependency, diminished quality of life, self-harming behaviors, and an elevated risk of suicide [5]. The prevalence of depression among university students in China has been reported to have increased from 33.6% to 35.4% between 2015 and 2018 [6]. Moreover, individuals were found to be more susceptible to depression during the COVID-19 pandemic compared to the pre-pandemic period [7]. Therefore, COVID-19 is expected to have a negative impact on the mental health of university students in China. The incidence of depression is higher in concentrated isolation under the COVID-19 epidemic, and the negative subjective perception of isolation and high stress level are significant risk factors for the occurrence of depression [8]. Given the impact of mental health on the overall well-being and development of contemporary university students, it is essential to collect data on depression prevalence among university students in China and implement necessary interventions [9].

Materials and Methods.

Objective: The purpose of this review is to investigate and analyze the prevalence of depression among university students in China. The findings of this study aim to provide a direction for psychological intervention and the improvement of depression among university students in China.

Methods: The protocol was developed based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocols (PRISMA-P) and registered in PROSPERO (CRD42024502949) [10].

Eligibility criteria: This study included cross-sectional studies that examined the prevalence of depression among university students in China. Depression, in this context, was defined as a depressed mood based on a scale that identified positive participants. The sample size of the studies did not exceed 300, and the original data was complete. The selected literature excluded studies that had poor quality, unavailable raw data, or focused on stress research in emergency or special situations (e.g., earthquakes, influenza). Additionally, literature that was not consistent with the theme (e.g., reviews, reports) or not in Chinese or English languages was also excluded.

Information sources: The searches were conducted in the following databases: PubMed, CNKI, Web of Science, and Wang-fang database.

© *GMN* 234

Search strategy: A combination of terms was searched: "depression" or "depressive disorder," "university" or "university students," "China" or "Chinese," and "cross-sectional study." No restrictions were applied to language, publication status, and publication time. The study screening process is clarified for systematic reviews and meta-analyses flow charts (Figure 1).

Figure 1 Flow diagram of study selection process.

Two reviewers worked independently to conduct title and abstract screening and full-text reviews using pre-defined eligibility criteria and standardized forms. Disagreements were resolved by a third reviewer.




Figure 1. HBOT Treatment table for Long COVID.

Results.

Data collection process: Reviewers conducted abstractions independently and in duplicate using standardized forms. Discrepancies were resolved by consensus, with input from a third reviewer if needed. For missing data, reviewers attempted to contact study authors when possible.

Data items:

The following study data was abstracted from each study:

- Study citation and author contact details.
- Study design, duration, and setting.
- Country.
- Number of participants.
- Basic information of participants (age, sex, major).
- Prevalence (male, female).

Outcomes and prioritization:

The primary outcome was the prevalence of depression among Chinese university students, while secondary outcomes are associated with risk or protective factors of depression.

Risk of bias in individual studies:

Standards that the Agency for Healthcare Research and Quality (AHRQ) recommended were used to evaluate the quality of observational studies. The criteria for evaluating a cross-sectional study consisted of 11 items answered with "yes," "no," and "unclear" [11].

Define the source of information (survey, record review).

List inclusion and exclusion criteria for exposed and unexposed subjects (cases and controls) or refer to previous publications.

Indicate the time period used for identifying patients.

Indicate whether or not subjects were consecutive if not population based.

Indicate if evaluators of subjective components of the study were masked to other aspects of the status of the participants.

Describe any assessments undertaken for quality assurance purposes (e.g., test/retest of primary outcome measurements).

Explain any patient exclusions from the analysis.

Describe how confounding was assessed and/or controlled.

If applicable, explain how missing data were handled in the analysis.

Summarize patient response rates and completeness of data collection.

Clarify what follow-up, if any, was expected and the percentage of patients for which incomplete data or follow-up was obtained [12].

Data synthesis and summary measures:

Statistical analysis was carried out by using STATA 11.0 software [13]. When $P \ge 0.1$ and $I^2 < 50\%$, there was no significant statistical heterogeneity, then a fixed effect model was adopted. P < 0.1 and $I^2 \ge 50\%$ suggested statistical heterogeneity, then the random-effects model was used for combined analysis [14]. Subgroup analysis was used to explore the source of heterogeneity. Publication bias was measured using funnel plots [13].

Subgroup analysis and sensitivity analysis:

Regardless of statistical heterogeneity, the subgroup analysis was conducted. Sex, region, and whether or not the study was carried out before COVID-19 were identified as factors for subgroup analysis. In addition, a one-by-one elimination method was adopted in sensitivity analysis.

Discussion.

Our systematic review aims to analyze the prevalence of depression among university students in China, contributing to the implementation of mental health education and psychological interventions for this population. By developing mental health education programs and prioritizing the mental well-being of university students, we can effectively address mental health issues such as depression and improve their academic performance and overall quality of life.

However, this review has potential limitations. Firstly, the definition of depression in this study is based on self-reported depressed mood, which may lead to an overestimation of prevalence rates. Additionally, the included studies are cross-sectional in nature, resulting in heterogeneity in study designs. To mitigate this, subgroup analyses were conducted to account for influential factors of depression.

Conclusion.

The findings from our meta-analysis will inform the development of mental health education initiatives for university students and community mental health efforts.

Moreover, further researches will be conducted to study current depression levels of depression among university students in

China to find out how did COVID-19 epidemic affect students' mental health.

Conflict-of-interest statement.

The authors declare that this research was conducted in the absence of any business or financial relationships that could be construed as potential conflicts of interest.

PRISMA 2009 Checklist statement.

The authors declare that this research was conducted in the absence of any business or financial relationships that could be construed as potential conflicts of interest.

Acknowledgements.

The research was supported by 2023 Taizhou University Higher Education Teaching Reform Project (No.79 and No.78), Science and Technology Association Soft Science Research Project of Taizhou City (No. KT202401100).

REFERENCES

- 1. Nobile M, Cataldo GM, Marino C, et al. Diagnosis and treatment of dysthymia in children and adolescents. CNS Drugs. 2003;17:927-946.
- 2. Rakel RE. Depression. Prim Care. 1999;26:211-224.
- 3. Satyanarayana PT, Suryanarayana R, Yesupatham ST, et al. Does Vitamin D3 Supplementation Improve Depression Scores among Rural Adolescents? A Randomized Controlled Trial. Nutrients. 2024;16.
- 4. Liu J, Kong J, Zhang X. Study on Differences between Patients with Physiological and Psychological Diseases in Online Health Communities: Topic Analysis and Sentiment Analysis. Int J Environ Res Public Health. 2020;17.
- 5. Ebert DD, Buntrock C, Mortier P, et al. Prediction of major depressive disorder onset in college students. Depress Anxiety. 2019;36:294-304.

- 6. Tang W, Dai Q. Depressive symptoms among first-year Chinese undergraduates: The roles of socio-demographics, coping style, and social support. Psychiatry Res. 2018;270:89-96.
- 7. Zhao C, He J, Du M, et al. A cross-lagged study of the associations among problematic smartphone use, depressive symptoms, and suicidal ideation in chinese university students. Curr Psychol. 2023:1-12.
- 8. Filiz B, Özyol FC, Güven B, et al. Physical activity, anxiety, depression, and coping in Turkish men and women during the first wave of COVID-19. Cad Saude Publica. 2024;40:e00057123.
- 9. Liu W, Zhang R, Wang H, et al. Association between anxiety, depression symptoms, and academic burnout among Chinese students: the mediating role of resilience and self-efficacy. BMC Psychol. 2024;12:335.
- 10. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. Syst Rev. 2015;4:1.
- 11. Zhang Y, Chen X, Lao Y, et al. Effects of the Implementation of Intelligent Technology for Hand Hygiene in Hospitals: Systematic Review and Meta-analysis. J Med Internet Res. 2023;25:e37249.
- 12. Lang S, Kleijnen J. Quality assessment tools for observational studies: lack of consensus. Int J Evid Based Healthc. 2010;8:247.
- 13. Song Y, Song N, Jia L, et al. Effectiveness and safety of Danshen injections in treatment of cardiac failure: a network meta-analysis. Front Pharmacol. 2024;15:1319551.
- 14. Dehkordi SR, Pahlavani N, Nikbaf-Shandiz M, et al. A systematic review, meta-analysis, dose-response, and meta-regression of the effects of acarbose intake on glycemic markers in adults. J Diabetes Metab Disord. 2024;23:135-172.