

# GEORGIAN MEDICAL NEWS

---

ISSN 1512-0112

NO 2 (359) Февраль 2025

---

ТБИЛИСИ - NEW YORK



ЕЖЕМЕСЯЧНЫЙ НАУЧНЫЙ ЖУРНАЛ

Медицинские новости Грузии  
საქართველოს სამედიცინო სიახლენი

## GEORGIAN MEDICAL NEWS

Monthly Georgia-US joint scientific journal published both in electronic and paper formats of the Agency of Medical Information of the Georgian Association of Business Press.  
Published since 1994. Distributed in NIS, EU and USA.

**GMN: Georgian Medical News** is peer-reviewed, published monthly journal committed to promoting the science and art of medicine and the betterment of public health, published by the GMN Editorial Board since 1994. GMN carries original scientific articles on medicine, biology and pharmacy, which are of experimental, theoretical and practical character; publishes original research, reviews, commentaries, editorials, essays, medical news, and correspondence in English and Russian.

GMN is indexed in MEDLINE, SCOPUS, PubMed and VINITI Russian Academy of Sciences. The full text content is available through EBSCO databases.

**GMN: Медицинские новости Грузии** - ежемесячный рецензируемый научный журнал, издаётся Редакционной коллегией с 1994 года на русском и английском языках в целях поддержки медицинской науки и улучшения здравоохранения. В журнале публикуются оригинальные научные статьи в области медицины, биологии и фармации, статьи обзорного характера, научные сообщения, новости медицины и здравоохранения. Журнал индексируется в MEDLINE, отражён в базе данных SCOPUS, PubMed и ВИНТИ РАН. Полнотекстовые статьи журнала доступны через БД EBSCO.

**GMN: Georgian Medical News** – საქართველოს სამედიცინო სიახლენი – არის ყოველთვიური სამეცნიერო სამედიცინო რეცენზირებადი ჟურნალი, გამოიცემა 1994 წლიდან, წარმოადგენს სარედაქციო კოლეგიისა და აშშ-ის მეცნიერების, განათლების, ინდუსტრიის, ხელოვნებისა და ბუნებისმეტყველების საერთაშორისო აკადემიის ერთობლივ გამოცემას. GMN-ში რუსულ და ინგლისურ ენებზე ქვეყნდება ექსპერიმენტული, თეორიული და პრაქტიკული ხასიათის ორიგინალური სამეცნიერო სტატიები მედიცინის, ბიოლოგიისა და ფარმაციის სფეროში, მიმოხილვითი ხასიათის სტატიები.

ჟურნალი ინდექსირებულია MEDLINE-ის საერთაშორისო სისტემაში, ასახულია SCOPUS-ის, PubMed-ის და ВИНТИ РАН-ის მონაცემთა ბაზებში. სტატიების სრული ტექსტი ხელმისაწვდომია EBSCO-ს მონაცემთა ბაზებიდან.

### WEBSITE

[www.geomednews.com](http://www.geomednews.com)

## К СВЕДЕНИЮ АВТОРОВ!

При направлении статьи в редакцию необходимо соблюдать следующие правила:

1. Статья должна быть представлена в двух экземплярах, на русском или английском языках, напечатанная через **полтора интервала на одной стороне стандартного листа с шириной левого поля в три сантиметра**. Используемый компьютерный шрифт для текста на русском и английском языках - **Times New Roman (Кириллица)**, для текста на грузинском языке следует использовать **AcadNusx**. Размер шрифта - **12**. К рукописи, напечатанной на компьютере, должен быть приложен CD со статьей.

2. Размер статьи должен быть не менее десяти и не более двадцати страниц машинописи, включая указатель литературы и резюме на английском, русском и грузинском языках.

3. В статье должны быть освещены актуальность данного материала, методы и результаты исследования и их обсуждение.

При представлении в печать научных экспериментальных работ авторы должны указывать вид и количество экспериментальных животных, применявшиеся методы обезболивания и усыпления (в ходе острых опытов).

4. К статье должны быть приложены краткое (на полстраницы) резюме на английском, русском и грузинском языках (включающее следующие разделы: цель исследования, материал и методы, результаты и заключение) и список ключевых слов (key words).

5. Таблицы необходимо представлять в печатной форме. Фотокопии не принимаются. **Все цифровые, итоговые и процентные данные в таблицах должны соответствовать таковым в тексте статьи**. Таблицы и графики должны быть озаглавлены.

6. Фотографии должны быть контрастными, фотокопии с рентгенограмм - в позитивном изображении. Рисунки, чертежи и диаграммы следует озаглавить, пронумеровать и вставить в соответствующее место текста **в tiff формате**.

В подписях к микрофотографиям следует указывать степень увеличения через окуляр или объектив и метод окраски или импрегнации срезов.

7. Фамилии отечественных авторов приводятся в оригинальной транскрипции.

8. При оформлении и направлении статей в журнал МНГ просим авторов соблюдать правила, изложенные в «Единых требованиях к рукописям, представляемым в биомедицинские журналы», принятых Международным комитетом редакторов медицинских журналов - <http://www.spinesurgery.ru/files/publish.pdf> и [http://www.nlm.nih.gov/bsd/uniform\\_requirements.html](http://www.nlm.nih.gov/bsd/uniform_requirements.html) В конце каждой оригинальной статьи приводится библиографический список. В список литературы включаются все материалы, на которые имеются ссылки в тексте. Список составляется в алфавитном порядке и нумеруется. Литературный источник приводится на языке оригинала. В списке литературы сначала приводятся работы, написанные знаками грузинского алфавита, затем кириллицей и латиницей. Ссылки на цитируемые работы в тексте статьи даются в квадратных скобках в виде номера, соответствующего номеру данной работы в списке литературы. Большинство цитированных источников должны быть за последние 5-7 лет.

9. Для получения права на публикацию статья должна иметь от руководителя работы или учреждения визу и сопроводительное отношение, написанные или напечатанные на бланке и заверенные подписью и печатью.

10. В конце статьи должны быть подписи всех авторов, полностью приведены их фамилии, имена и отчества, указаны служебный и домашний номера телефонов и адреса или иные координаты. Количество авторов (соавторов) не должно превышать пяти человек.

11. Редакция оставляет за собой право сокращать и исправлять статьи. Корректур авторам не высылаются, вся работа и сверка проводится по авторскому оригиналу.

12. Недопустимо направление в редакцию работ, представленных к печати в иных издательствах или опубликованных в других изданиях.

**При нарушении указанных правил статьи не рассматриваются.**

## REQUIREMENTS

Please note, materials submitted to the Editorial Office Staff are supposed to meet the following requirements:

1. Articles must be provided with a double copy, in English or Russian languages and typed or computer-printed on a single side of standard typing paper, with the left margin of 3 centimeters width, and 1.5 spacing between the lines, typeface - **Times New Roman (Cyrillic)**, print size - 12 (referring to Georgian and Russian materials). With computer-printed texts please enclose a CD carrying the same file titled with Latin symbols.

2. Size of the article, including index and resume in English, Russian and Georgian languages must be at least 10 pages and not exceed the limit of 20 pages of typed or computer-printed text.

3. Submitted material must include a coverage of a topical subject, research methods, results, and review.

Authors of the scientific-research works must indicate the number of experimental biological species drawn in, list the employed methods of anesthetization and soporific means used during acute tests.

4. Articles must have a short (half page) abstract in English, Russian and Georgian (including the following sections: aim of study, material and methods, results and conclusions) and a list of key words.

5. Tables must be presented in an original typed or computer-printed form, instead of a photocopied version. **Numbers, totals, percentile data on the tables must coincide with those in the texts of the articles.** Tables and graphs must be headed.

6. Photographs are required to be contrasted and must be submitted with doubles. Please number each photograph with a pencil on its back, indicate author's name, title of the article (short version), and mark out its top and bottom parts. Drawings must be accurate, drafts and diagrams drawn in Indian ink (or black ink). Photocopies of the X-ray photographs must be presented in a positive image in **tiff format**.

Accurately numbered subtitles for each illustration must be listed on a separate sheet of paper. In the subtitles for the microphotographs please indicate the ocular and objective lens magnification power, method of coloring or impregnation of the microscopic sections (preparations).

7. Please indicate last names, first and middle initials of the native authors, present names and initials of the foreign authors in the transcription of the original language, enclose in parenthesis corresponding number under which the author is listed in the reference materials.

8. Please follow guidance offered to authors by The International Committee of Medical Journal Editors guidance in its Uniform Requirements for Manuscripts Submitted to Biomedical Journals publication available online at: [http://www.nlm.nih.gov/bsd/uniform\\_requirements.html](http://www.nlm.nih.gov/bsd/uniform_requirements.html)  
[http://www.icmje.org/urm\\_full.pdf](http://www.icmje.org/urm_full.pdf)

In GMN style for each work cited in the text, a bibliographic reference is given, and this is located at the end of the article under the title "References". All references cited in the text must be listed. The list of references should be arranged alphabetically and then numbered. References are numbered in the text [numbers in square brackets] and in the reference list and numbers are repeated throughout the text as needed. The bibliographic description is given in the language of publication (citations in Georgian script are followed by Cyrillic and Latin).

9. To obtain the rights of publication articles must be accompanied by a visa from the project instructor or the establishment, where the work has been performed, and a reference letter, both written or typed on a special signed form, certified by a stamp or a seal.

10. Articles must be signed by all of the authors at the end, and they must be provided with a list of full names, office and home phone numbers and addresses or other non-office locations where the authors could be reached. The number of the authors (co-authors) must not exceed the limit of 5 people.

11. Editorial Staff reserves the rights to cut down in size and correct the articles. Proof-sheets are not sent out to the authors. The entire editorial and collation work is performed according to the author's original text.

12. Sending in the works that have already been assigned to the press by other Editorial Staffs or have been printed by other publishers is not permissible.

**Articles that Fail to Meet the Aforementioned  
Requirements are not Assigned to be Reviewed.**

## ავტორთა საქურაღებოლ!

რედაქციაში სტატიის წარმოდგენისას საჭიროა დაიცვათ შემდეგი წესები:

1. სტატია უნდა წარმოადგინოთ 2 ცალად, რუსულ ან ინგლისურ ენებზე დაბეჭდილი სტანდარტული ფურცლის 1 გვერდზე, 3 სმ სიგანის მარცხენა ველისა და სტრიქონებს შორის 1,5 ინტერვალის დაცვით. გამოყენებული კომპიუტერული შრიფტი რუსულ და ინგლისურენოვან ტექსტებში - **Times New Roman (Кириллица)**, ხოლო ქართულენოვან ტექსტში საჭიროა გამოვიყენოთ **AcadNusx**. შრიფტის ზომა – 12. სტატიას თან უნდა ახლდეს CD სტატიით.

2. სტატიის მოცულობა არ უნდა შეადგენდეს 10 გვერდზე ნაკლებს და 20 გვერდზე მეტს ლიტერატურის სიის და რეზიუმეების (ინგლისურ, რუსულ და ქართულ ენებზე) ჩათვლით.

3. სტატიაში საჭიროა გაშუქდეს: საკითხის აქტუალობა; კვლევის მიზანი; საკვლევი მასალა და გამოყენებული მეთოდები; მიღებული შედეგები და მათი განსჯა. ექსპერიმენტული ხასიათის სტატიების წარმოდგენისას ავტორებმა უნდა მიუთითონ საექსპერიმენტო ცხოველების სახეობა და რაოდენობა; გაუტკივარებისა და დაძინების მეთოდები (მწვავე ცდების პირობებში).

4. სტატიას თან უნდა ახლდეს რეზიუმე ინგლისურ, რუსულ და ქართულ ენებზე არანაკლებ ნახევარი გვერდის მოცულობისა (სათაურის, ავტორების, დაწესებულების მითითებით და უნდა შეიცავდეს შემდეგ განყოფილებებს: მიზანი, მასალა და მეთოდები, შედეგები და დასკვნები; ტექსტუალური ნაწილი არ უნდა იყოს 15 სტრიქონზე ნაკლები) და საკვანძო სიტყვების ჩამონათვალი (key words).

5. ცხრილები საჭიროა წარმოადგინოთ ნაბეჭდი სახით. ყველა ციფრული, შემაჯამებელი და პროცენტული მონაცემები უნდა შეესაბამებოდეს ტექსტში მოყვანილს.

6. ფოტოსურათები უნდა იყოს კონტრასტული; სურათები, ნახაზები, დიაგრამები - დასათაურებული, დანომრილი და სათანადო ადგილას ჩასმული. რენტგენოგრაფიების ფოტოასლები წარმოადგინეთ პოზიტიური გამოსახულებით **tiff** ფორმატში. მიკროფოტოსურათების წარწერებში საჭიროა მიუთითოთ ოკულარის ან ობიექტივის საშუალებით გადიდების ხარისხი, ანათალებების შედეგების ან იმპრეგნაციის მეთოდი და აღნიშნოთ სურათის ზედა და ქვედა ნაწილები.

7. სამამულო ავტორების გვარები სტატიაში აღინიშნება ინიციალების თანდართვით, უცხოურისა – უცხოური ტრანსკრიპციით.

8. სტატიას თან უნდა ახლდეს ავტორის მიერ გამოყენებული სამამულო და უცხოური შრომების ბიბლიოგრაფიული სია (ბოლო 5-8 წლის სიღრმით). ანბანური წყობით წარმოდგენილ ბიბლიოგრაფიულ სიაში მიუთითეთ ჯერ სამამულო, შემდეგ უცხოელი ავტორები (გვარი, ინიციალები, სტატიის სათაური, ჟურნალის დასახელება, გამოცემის ადგილი, წელი, ჟურნალის №, პირველი და ბოლო გვერდები). მონოგრაფიის შემთხვევაში მიუთითეთ გამოცემის წელი, ადგილი და გვერდების საერთო რაოდენობა. ტექსტში კვადრატულ ფხიხლებში უნდა მიუთითოთ ავტორის შესაბამისი N ლიტერატურის სიის მიხედვით. მიზანშეწონილია, რომ ციტირებული წყაროების უმეტესი ნაწილი იყოს 5-6 წლის სიღრმის.

9. სტატიას თან უნდა ახლდეს: ა) დაწესებულების ან სამეცნიერო ხელმძღვანელის წარდგინება, დამოწმებული ხელმოწერითა და ბეჭდით; ბ) დარგის სპეციალისტის დამოწმებული რეცენზია, რომელშიც მითითებული იქნება საკითხის აქტუალობა, მასალის საკმაობა, მეთოდის სანდოობა, შედეგების სამეცნიერო-პრაქტიკული მნიშვნელობა.

10. სტატიის ბოლოს საჭიროა ყველა ავტორის ხელმოწერა, რომელთა რაოდენობა არ უნდა აღემატებოდეს 5-ს.

11. რედაქცია იტოვებს უფლებას შეასწოროს სტატია. ტექსტზე მუშაობა და შეჯერება ხდება საავტორო ორიგინალის მიხედვით.

12. დაუშვებელია რედაქციაში ისეთი სტატიის წარდგენა, რომელიც დასაბეჭდად წარდგენილი იყო სხვა რედაქციაში ან გამოქვეყნებული იყო სხვა გამოცემებში.

აღნიშნული წესების დარღვევის შემთხვევაში სტატიები არ განიხილება.

Gavrilova Uliana V, Alieva Samaya A, Gerasimenko Aleksandra A, Mikhaleva Ekaterina A, Solovieva Ekaterina V, Vedzizheva Khava Kh, Sadykov Magomed L, Belousova Anastasia A, Ladaev Abubakar Kh, Aupov Ibragim A, Maasheva Elita A, Makhamaev Ilias B, Yanarcae V Ibragim R. NEXT-GENERATION ANTIOXIDANTS: SHOULD WE TARGET PEROXIREDOXINS (PRX)?.....	6-14
Marina Endeladze, Maia Zhamutashvili, Tinatin Gognadze, Elene Meskhi, Natia Jojua, M. Akhvlediani. CASE REPORT OF CAT SCRATCH DISEASE (BARTONELLA).....	15-17
Karapetyan A.G, Santini C, Pellei M, Caviglia M, Dallakyan A.M, Petrosyan Zh.H, Danielyan M.H, Nebogova K.A, Grigoryan V.S. MANIFESTATION OF RADIOPROTECTIVE PROPERTIES IN COPPER COMPLEXES [CU(L <sup>CE3</sup> ) <sub>2</sub> ] AND [CU(ADM)(PPH <sub>3</sub> ) <sub>2</sub> ]PF <sub>6</sub> .....	18-22
Nato Nakudashvili, Levan Ratiani, Tamar Megrelishvili, Elene Saribekovi, Marine Tsabadze, Nina Kipiani, Nino Intskirveli, Magda Tortladze, Tea Gabunia, Shorena Tsiklauri, Zaza Nakudashvili, George Ormotsadzr, Tamar Sanikidze. FEATURES OF VASOMOTOR RHINITIS (VMR) IN PATIENTS WITH A HISTORY OF COVID-19 INFECTION.....	23-31
Warda Mohamed, Rashid Eltayeb, Hussam Ali Osman, Mosab Omer, Elryah. I. Ali, Ammar Abdelmola, Tagwa Yousif, Mohamed Belhocine, Safa Taha, Elyasa Elfaki, Wael Alzahrani, Asaad Babker, Abdelgadir Elamin Eltom, Marwan Ismail, Ayman Alfeel. BIOCHEMICAL INSIGHTS AND HORMONAL MARKERS OF POLYCYSTIC OVARY SYNDROME IN SUDANESE WOMEN: EXPLORING INFERTILITY AND RISK FACTORS IN RED SEA STATE.....	32-39
Kolupayev S.M, Goloborodko M.M, Bytiak S.Yu, Lavrinenko A.S, Lupyr M, Lantukh I.V, Lytvynova I.L, Gulbs O.A, Dikhtyarenko S.Yu, Kobets O.V. PSYCHOLOGICAL FEATURES OF THE REHABILITATION OF PERSONS WITH UROLITHIASIS.....	40-44
Wang-Yong Li, Xiao-Yan Yang, Zhun Cai, Guo-Fu Chen. A CASE OF CECAL CANCER WITH RETROPERITONEAL ABSCESS.....	45-48
Tamriko Dzotsenidze, Arsen Gvenetadze, Giorgi Burkadze, Ekaterine Isakidi, Mariam Shulaia. GENETIC ALTERATIONS IN TUBO-OVARIAN EPITHELIUM DURING OVARIAN NEOPLASIA.....	49-54
Ali M. Muhammed Ali, Omar M. Yahya, Ehsan HT. AlDabbagh. IN SILICO DOCKING OF Silymarin ACTIVE CONSTITUENTS WITH INSULIN RECEPTORS: A STEP TOWARD DIABETES THERAPEUTICS.....	55-62
Armenuhi Avagyan, Taline K. Ashekian, Armenuhi Snkhchyan, Hasmik Mkrtychyan, Tigran Petrosyan. BARRIERS AND SOLUTIONS IN THE USE OF ALTERNATIVE AND AUGMENTATIVE COMMUNICATION: A PILOT DESCRIPTIVE STUDY PRESENTING INSIGHTS FROM ARMENIAN PROFESSIONALS.....	63-71
Xin-Juan Wang, Lian-Ping He. STATISTICAL TEACHING ON BUILDING STROKE PREDICTION MODELS.....	72-75
Kosherova Bakhyt Nurgaliyevna, Abbozova Shakhnosa Maratovna, Smagul Manar Asyrovna, Zhumagaliyeva Galina Dautovna, Sagyndykova Togzhan Baibolsynovna. MEASLES IN PREGNANCY IN THE REPUBLIC OF KAZAKHSTAN: CLINICAL AND LABORATORY MANIFESTATIONS AND OUTCOMES.....	76-80
Warda Mohamed, Rashid Eltayeb, Hussam Ali Osman, Mosab Omer, Elryah. I. Ali, Ammar Abdelmola, Tagwa Yousif, Mohamed Belhocine, Safa Taha, Elyasa Elfaki, Wael Alzahrani, Ayman Alfeel, Asaad Babker, Abdelgadir Elamin Eltom, Marwan Ismail. FOLLICLE-STIMULATING HORMONE RECEPTOR MUTATIONS IN SUDANESE WOMEN: A STUDY ON POLYCYSTIC OVARY SYNDROME.....	81-86
Saidulaev M.A, Osipova N.I, Gurtskaya A.D, Semov N.D, Khusainov R.R, Khabarov E.A, Lech D.S, Stotland P.A, Eloeva V.V, Syromyatnikova A.V. PIRACETAM HELPS RESTORE VISION AFTER CRANIOCEREBRAL TRAUMA.....	87-88
Ming Li, Wen-Wen Hao, Li-Juan Ru. THE ASSOCIATION BETWEEN AQUEOUS HUMOR MICROENVIRONMENT IN DIABETIC CATARACT PATIENTS AND POSTOPERATIVE MACULAR EDEMA AND VISUAL ACUITY CHANGES.....	89-90
Voloshyn-Haponov I.K, Lantukh I.V, Gulbs O.A, Dikhtyarenko S.Yu, Kobets O.V, Pustova N.O, Popova N.G, Gridneva O.V, Ostapenko V.M, Mikhhanovska N.G, Torianyik I.I. PSYCHOLOGICAL FEATURES OF THE SUBJECTIVE PERCEPTION OF THE QUALITY OF LIFE OF PATIENTS WITH HEPATOCEREBRALDYSTROPHY.....	91-95
Bangqiang Hou, Wei Liu, Ke Pan, Yiya Wang, Yaomin Luo, Yutong Han, Jingjing Liu, Qing Wu, Yin Xu Wang. BRAIN NETWORK FUNCTIONAL CONNECTIVITY AND CORTICAL ACTIVATION FEATURES DURING THE SWALLOWING TASK FOR THE PATIENTS OF POST STROKE DYSPHAGIA: A MULTI- CHANNEL FNIRS STUDY.....	96-107
Assel Ibrayeva, Dinara Ospanova, Korlan Saduakasova, Anar Akshalova, Anar Muratbayeva, Shynar Tanabayeva, Ildar Fakhradiyev. PREVALENCE AND SOCIO-DEMOGRAPHIC RISK FACTORS OF EMOTIONAL BURNOUT AMONG PSYCHIATRISTS AND NURSING STAFF IN PSYCHIATRIC SERVICES IN KAZAKHSTAN.....	108-115

Violeta Grajčevci Uka, Art Uka, Lirim Isufi. THE SOCIODEMOGRAPHICAL AND MORPHOLOGICAL CHARACTERISTICS OF PRESCHOOL CHILDREN WITH SIDEROPENIC ANEMIA IN THE KOSOVO SAMPLE.....	116-119
Li-Bo Wang, Chun-Miao Xu. ENHANCING OPHTHALMIC NURSING EDUCATION: A COMPREHENSIVE APPROACH TO CLINICAL TEACHING AND TRAINING.....	120-122
Maia Zhamutashvili, Tinatin Gognadze, Natia Jojua, Elene Meskhi, Ketevan Meskhi, Ekaterine Dolmazashvili. CO-OCCURRENCE OF HANTAVIRUS PULMONARY SYNDROME AND HEMORRHAGIC FEVER WITH RENAL SYNDROME.....	123-125
Abdualikova D.B, Auezova A.M, Baymuratova M.A, Yessembayeva S.S, Yermukhanova G.T, Yerkibayeva ZH.U. WORKSHOP AS A PRACTICE-ORIENTED METHOD OF TEACHING DENTISTS: INTRODUCTION TO EDUCATIONAL PROGRAMS FOR WORKING WITH CHILDREN WITH AUTISM SPECTRUM DISORDERS (ASD).....	126-132
Li-Juan Ru, Qian-Qian Yao, Ming Li. RISK PREDICTION MODEL FOR ACUTE KIDNEY INJURY IN PATIENTS WITH SEVERE ACUTE PANCREATITIS.....	133-135
Sawer S. Ahmed, Abdulazeez M. Brifkani, Haval J. Ali, Nasir A. Al Allawi. FACTORS AFFECTING HEALTH RELATED QUALITY OF LIFE IN ADULT PATIENTS WITH BETA-THALASSEMIA MAJOR.....	136-142
Liu-Xia Shi, Xiao-Ya Peng, Xiao-Xu Ruan, Rui Li, Wen-Jie Wen, Chao Deng. WHOLE TRANSCRIPTOME SEQUENCING AND CIRC_HSA_0001847 ON PROLIFERATION AND INVASION OF ORAL SQUAMOUS CELL CARCINOMA.....	143-155
Takuma Hayashi, Krishna Prasad Acharya, Sarita Phuyal, Ikuo Konishi. THE ROLE OF LIVE BIRD MARKETS SHOULD BE EMPHASIZED IN PREVENTING THE SPREAD OF HIGHLY PATHOGENIC AVIAN INFLUENZA INFECTIONS.....	156-158
Karapetyan A.G, Grigoryan V.S, Santini C, Pellei M, Del Gobbo J, Dallakyan A.M, Petrosyan Zh.H, Fanarjyan S.A, Danielyan M.H, Nebogova K.A. BLOOD AND CYTOGENETIC MARKERS IN EXPERIMENTAL BURNS AND THEIR TREATMENT WITH [CU(L <sup>CF3</sup> ) <sub>2</sub> ] AND [CU(ADM)(PPH <sub>3</sub> ) <sub>2</sub> ]PF <sub>6</sub> .....	159-163
Sharofova M.U, Khalimova F.T, Habasi Maidina, Jiangyu Zhao, Haji Akber Aisa. ANTIMICROBIAL AND ANTI-INFLAMMATORY ACTIVITY OF PLANT EXTRACTS: PROSPECTS FOR THE DEVELOPMENT OF COMBINED THERAPEUTIC AGENTS.....	164-168

## BIOCHEMICAL INSIGHTS AND HORMONAL MARKERS OF POLYCYSTIC OVARY SYNDROME IN SUDANESE WOMEN: EXPLORING INFERTILITY AND RISK FACTORS IN RED SEA STATE

Warda Mohamed<sup>1</sup>, Rashid Eltayeb<sup>2</sup>, Hussam Ali Osman<sup>3</sup>, Mosab Omer<sup>1</sup>, Elryah. I. Ali<sup>4</sup>, Ammar Abdelmola<sup>5</sup>, Tagwa Yousif<sup>5</sup>, Mohamed Belhocine<sup>6</sup>, Safa M. Taha<sup>7</sup>, Wael Alzahrani<sup>8</sup>, Elyasa Elfaki<sup>8</sup>, Asaad Babker<sup>9</sup>, Abdelgadir Elamin Eltom<sup>9</sup>, Marwan Ismail<sup>9</sup>, Ayman Alfeel<sup>8\*</sup>.

<sup>1</sup>Faculty of Medical Laboratory Sciences, Clinical Chemistry Department, Shendi university

<sup>2</sup>Department of Biochemistry, Faculty of Medicine- Shendi University-Sudan.

<sup>3</sup>Faculty of Medical and Health Sciences, Liwa College, Abu Dhabi, UAE.

<sup>4</sup>Department of Medical Laboratory Technology, College of Applied Medical Sciences, Northern Border University, Arar, Saudi Arabia.

<sup>5</sup>Department of Medical Laboratory Technology, College of Nursing and Health Sciences, Jazan University, Jazan, 45142, Saudi Arabia.

<sup>6</sup>Department of Molecular Medicine, Al-Jawhara Centre for Molecular Medicine, Genetics, and Inherited Disorders, Arabian Gulf University, Manama, Bahrain.

<sup>7</sup>Arabian Gulf University, Molecular Medicine department, Kingdom of Bahrain.

<sup>8</sup>Department of Clinical Laboratory Science, College of Applied Medical Sciences-Qurayyat, Jouf University, Al Jouf, Saudi Arabia.

<sup>9</sup>Department of Medical Laboratory Sciences, College of Health Sciences, Gulf Medical University.

### Abstract.

**Introduction:** Polycystic Ovary Syndrome (PCOS) is a prevalent hormonal condition affecting women of reproductive age, with prevalence rates ranging from 5% to 13%. Symptoms include irregular menstrual cycles, androgen excess, hair growth, acne, and cysts in the ovaries, often seen during adolescence. **Aim:** The purpose of this study was to investigate the hormonal and biochemical profiles of women with PCOS in the Red Sea State, Sudan with a focus on infertility and associated risk factors. **Materials and methods:** This cross-sectional study involved 40 women with a Rotterdam-defined PCOS diagnosis and 40 healthy control subjects. Participants provided demographic information, including age, family history, irregular menstruation, infertility, metabolic abnormalities, and clinical symptoms. Serum was extracted to estimate hormonal profiles, and enzymatic techniques were used to analyze fasting blood glucose and lipid profiles, while ELISA quantitative methods were used for hormone profiles. **Results:** The study found that 40% of patients were aged 20-30, 45% were with primary schools' education level, 53% had low economic status, 58% had abnormal menstrual cycle features, 60% had a positive case history, 75% did not have chronic disease characteristics, 68% had infants, and 83% had normal body mass index. The study found an insignificant correlation between infertility hormones and lipid profiles in patients, with FSH, LH, and LDL showing insignificant variation, while PRL, Chol, TG, and HDL showed significant correlations. **Conclusion:** The study reveals significant changes in lipid and hormonal profiles in Sudanese women with PCOS, even those with normal BMI. Although cyst laterality doesn't strongly correlate with clinical indicators, more research is needed to understand its relationship with LH and chronic illness. Early screening and multidisciplinary therapy are crucial due to high incidence of dyslipidemia, positive family history, and irregular menstruation.

**Key words.** ELISA, Polycystic Ovary Syndrome (PCOS).

### Introduction.

Polycystic ovarian syndrome (PCOS), first described in 1935, is the most common endocrine pathology in female reproductive health, with prevalence ranging from 5% to 15%, diagnosed by chronic anovulation, hyperandrogenism, and polycystic ovaries [1].

According to a meta-analysis and systematic review, the overall prevalence of infertility in Sudan is roughly 13%, with 35% of cases being secondary infertility and 65% being primary infertility. Ovulatory abnormalities are common in infertile women, and PCOS accounts for 38% of these cases. Various factors lead to an increased prevalence of the disease, unknown prevalence on Sudan besides different diagnostic criteria used for diagnosis as well as the absence of screening program and genetic counseling in addition to the lack of awareness through the community about the importance of women health and its impact on her social life [2].

The typical biochemical abnormalities include increased levels of serum androgens (especially testosterone and androstenedione) and elevated Luteinizing Hormone (LH) levels, while Follicle-Stimulating Hormone (FSH) levels remain normal or are low [3].

Based on the "Rotterdam criteria," PCOS is diagnosed when two of the three characteristics—polycystic ovaries, anovulation, and androgen excess (clinical and/or biochemical)—are present [4]. The 2023 International PCOS Guideline criteria revise Rotterdam criteria, recommending diagnosis in adults based on oligo/amenorrhea, clinical/biochemical hyperandrogenism, and PCOM, and adolescents requiring both [5,6].

Many of the endocrine characteristics of PCOS seem to be caused by insulin resistance and its subsequent hyperinsulinemia in a significant number of these patients. Patients with PCOS have a five to ten times higher chance of developing type 2 diabetes mellitus than the general population [7].

PCOS is also a major contributor to infertility in women. For women with PCOS, improvements in lifestyle, such as decreasing weight, can lead to physical changes that make conception easier [8].



PCOS, including obesity, hirsutism, hair loss, irregular menstruation, and acne, can lead to low self-esteem, social rejection, and psychological problems. It also increases anxiety, depression, and loneliness, contributing to higher divorce rates and strained marriages.

Cultural, environmental, and genetic factors in Sudan may influence PCOS presentation and progression. Limited research on biochemical and hormonal underpinnings is crucial for effective diagnosis, management, and risk mitigation.

The purpose of this study is to investigate the hormonal and biochemical profiles of women with PCOS in the Red Sea State, with a focus on infertility and associated risk factors. By detecting patterns and connections, the study hopes to reveal insights that can improve diagnostic precision and inform targeted therapies. Furthermore, it investigates the impact of local environmental and lifestyle factors in regulating the clinical symptoms of PCOS, contributing to a better understanding of the disease in a geographically and culturally diverse population.

## Materials and Methods.

This an analytical laboratory-based cross-sectional study to investigate the hormonal and biochemical profiles of women with PCOS in the Red Sea State, with a focus on infertility and associated risk factors.

This cross-sectional study involved 40 women with a Rotterdam-defined PCOS diagnosis and 40 healthy control subjects. Participants provided demographic information, including age, family history, irregular menstruation, infertility, metabolic abnormalities, and clinical symptoms.

Any Sudanese females below the reproductive age, females with family history of adrenal hyperplasia, hyperprolactinemia, Cushing syndrome, thyroid disorders, ovarian cancer or breast cancer and causes of infertility rather than PCOS were excluded from the study.

Structured self-administered questionnaires were answered by study participants covering their; socio-demographic variables: Age, family history, menstrual abnormalities, infertility problems, metabolic abnormalities and clinical symptoms.

Venous blood was drawn from patients and controls with a disposable sterile plastic syringe. After a 10-minute centrifugation at 5000 RPM at room temperature, serum is recovered and stored at  $-80^{\circ}\text{C}$ .

Data was input and structured into a Microsoft Office Excel (2019) data sheet before being transferred to the statistical package program IBM SPSS Statistics for Windows, Version 26.049 to perform various statistical tests such as mean standard deviation and P-value.

For hormonal profile estimations, we used quantitative (ELISA) and enzymatic methods to analyze fasting blood glucose and lipid profile.

Shendi University's ethical committee approved the study, and participants provided their consent.

## Results.

This analytical, laboratory-based cross-sectional study was conducted to investigate various biochemical parameters in 40 Sudanese individuals with one or more female family members diagnosed with PCOS, alongside 40 individuals in a control

group. Descriptive and biological data were collected using a self-administered questionnaire. The information was analysed using SPSS version 26, with descriptive statistics applied to assess the frequency distributions of age (years), BMI, clinical signs and symptoms, and biochemical markers.

For both categories, the largest percentage of participants are in the 20–30 age range (Figure 1).

Compared to the control group, PCOS patients are more likely to have completed primary school. Both groups have comparable levels of secondary education. Although there are fewer women in both groups with a university degree, the case group's percentage is marginally greater (Figure 2).

The majority of women in both groups are low-income, the prevalence of PCOS is almost the same in the moderate-income group as it is in the controls, and there are more PCOS patients in the high-income group than in the controls (Figure 3).

Compared to healthy controls, women with PCOS have a higher prevalence of chronic illnesses, especially diabetes and hypertension (Figure 4)

75% of PCOS patients had abnormal menstruation compared to 67.5% in the control group Confirms that menstrual irregularity is more common in PCOS (Table 2), 60% of the PCOS group had a positive family history, suggesting a hereditary component in PCOS development (Table 3).

Only 50% of PCOS patients had regular medical follow-up indicates gaps in healthcare access or awareness (Table 3), 53% had bilateral cysts, 47% had unilateral suggest that bilateral involvement is slightly more common (Table 4) 68% of PCOS patients had a history of childbirth Implies that fertility is possible, though PCOS is a known infertility risk (Table 5).

83% had normal BMI, 17% were overweight Most PCOS patients in this sample were non-obese, highlighting that PCOS isn't limited to obese women (Table 6)

No significant associations found between cyst location and menstruation ( $p=0.2$ ), family history ( $p=0.5$ ), medical follow-up ( $p=0.1$ ) and history of delivery ( $p=0.5$ ) while Significant association found with: Chronic disease ( $p=0.04$ ), especially diabetes, which was more common in those with bilateral cysts (Tables 7-11)

Significant differences in PRL, Cholesterol, Triglycerides, and HDL (all  $p < 0.01$ ) – all higher in PCOS group while no significant differences in FSH, LH, and LDL (Table 12).

Significant difference in LH levels ( $p = 0.01$ ) higher in those with unilateral cysts however no significant differences in other hormones or lipids by cyst location (Table 13).

No statistically significant differences between regular and irregular menstruation in hormonal or lipid values with some trends (e.g., higher PRL in regular cycles), but not significant (Table 14).

No significant differences between normal and overweight groups, Overweight group had higher TG and lower HDL, but not statistically significant (Table 15).

## Discussion.

This analytical, laboratory-based cross-sectional study aimed to assess biochemical and clinical parameters associated with polycystic ovary syndrome (PCOS) among Sudanese women with familial PCOS history, compared to a control group. The




Figure 1. Age Distribution in Case and Control Groups.




Figure 2. Education Level Distribution in Case and Control Groups.




Figure 3. Economic Status Distribution in Case and Control Groups.




Figure 4. Chronic Disease Distribution in Case and Control Groups.

Table 1. Menstruation status patients and control under study.

Menstruation status		Frequency	Percent
Normal	Case	10	25%
	Control	13	32.5%
Abnormal	Case	30	75%
	Control	27	67.5%
Total		80	100%

Table 2. Family history in case group.

Family history	Frequency	Percent
Positive FH	24	60%
Negative FH	16	40%
Total	40	100%

Table 3. Medical follow up in case group.

Follow up	Frequency	Percent
Positive follow up	20	50%
Negative follow up	20	50%
Total	40	100%

Table 4. Cyst location in case group.

Cyst location	Frequency	Percent
Unilateral	19	47%
Bilateral	21	53%
Total	40	100%

Table 5. History of delivery in case group.

Patient has delivered before	Frequency	Percent
Yes	27	68%
No	13	32%
Total	40	100%

Table 6. BMI in case group.

BMI	Frequency	Percent
Normal	33	83%
Overweight	7	17%
Total	40	100%

Table 7. Association between cyst location and menstruation cycle status.

Cycle status		Cyst		Total	P.value
		Uni	Bi		
Normal	Count	7	15%	4	0.2
	%	17.5%	37.5%	10.0%	
Abnormal	Count	12%	6	4	
	%	30%	15%	10.0%	
Total	Count	19	21	40	
	%	100.0%	100.0%	100.0%	

The mean difference is significant at the (0.05) level.

Table 8. Association between cyst location and family history in patients.

Case history		Cyst		Total	P.value
		Uni	Bi		
Positive	Count	11	13	24	0.5
	%	27.5%	32.5%	60.0%	
Negative	Count	8	8	16	
	%	20	20%	40.0%	
Total	Count	19	21	40	
	%	100.0%	100.0%	100.0%	

\*The mean difference is significant at the (0.05) level.

Table 9. Relation between cyst location & medical follow up in patients.

Management		Cyst		Total	P.value
		Uni	Bi		
Positive	Count	7	13	20	0.1
	%	17.5%	32.5%	50.0%	
Negative	Count	12	8	20	
	%	30%	20%	50.0%	
Total	Count	19	21	40	
	%	100.0%	100.0%	100.0%	

\*The mean difference is significant at the (0.05) level.

Table 10. Association between cyst location and chronic diseases in patients.

Disease		Cyst		Total	P.value
		Uni	Bi		
DM	Count	0	6	6	0.04
	%	0.0%	15%	15.0%	
Hypertension	Count	2	2	4	
	%	5%	2%	10.0%	
No Disease	Count	17	13	30	
	%	42.5%	32.5%	75.0%	
Total	Count	19	21	40	
	%	100.0%	100.0%	100.0%	

\*The mean difference is significant at the (0.05) level.

Table 11. Association between cyst location & and delivery in patients.

Infant		Cyst		Total	P.value
		Uni	Bi		
Yes	Count	13	14	27	0.5
	%	32.5%	35%	67.5%	
No	Count	6	7	13	
	%	15%	17.5%	32.5%	
Total	Count	19	21	40	
	%	100.0%	100.0%	100.0%	

\*The mean difference is significant at the (0.05) level.

Table 12. Correlation between infertility hormones and lipid profile (case & control) in patients.

Sample		N	Mean	Std. Deviation	P.value
FSH (ng/dl)	Case	40	5.6	3.0	0.3
	Control	40	5.0	2.3	
LH (ng/dl)	Case	40	8.1	2.8	0.3
	Control	40	7.5	2.5	
PRL (ng/dl)	Case	40	31.2	12.8	0.00
	Control	40	14.7	5.5	
Chol (mmol/l)	Case	40	9.0	1.3	0.00
	Control	40	5.0	0.3	
TG (mmol/l)	Case	40	7.6	2.2	0.00
	Control	40	3.3	1.8	
HDL (mmol/l)	Case	40	4.0	1.7	0.00
	Control	40	1.8	0.3	
LDL (mmol/l)	Case	40	5.0	2.0	0.7
	Control	40	4.7	1.8	

\*The mean difference is significant at the (0.05) level.

Table 13. Correlation between infertility hormones & lipid profile with cyst location in patients.

Cyst		N	Mean	Std. Deviation	P.value
FSH (ng/dl)	Uni	19	3.9	1.6	0.08
	Bi	21	6.0	3.6	
LH (ng/dl)	Uni	19	9.0	2.1	0.01
	Bi	21	6.2	2.7	
PRL (ng/dl)	Uni	19	30.7	13.3	0.9
	Bi	21	31.5	12.7	
CHOL (mmol/l)	Uni	19	5.2	1.4	0.5
	Bi	21	5.5	1.3	
TG (mmol/l)	Uni	19	2.5	1.8	0.1
	Bi	21	4.1	2.3	
HDL (mmol/l)	Uni	19	3.8	1.6	0.7
	Bi	21	4.2	1.7	
LDL (mmol/l)	Uni	19	4.6	2.1	0.4
	Bi	21	5.0	2.0	

\*The mean difference is significant at the (0.05) level.

Table 14. Correlation between infertility hormones & lipid profile with menstruation in patients.

cycle status		N	Mean	Std. Deviation	P.value
FSH (ng/dl)	Irregular	23	4.7	2.6	0.6
	Regular	9	4.1	1.3	
LH (ng/dl)	Irregular	23	7.0	3.0	0.1
	Regular	9	9.1	1.3	
PRL (ng/dl)	Irregular	23	27.1	13.0	0.09
	Regular	9	39.4	8.1	
CHOL (mmol/l)	irregular	23	5.2	1.3	0.6
	Regular	9	5.6	1.4	
TG (mmol/l)	irregular	23	3.9	2.6	0.3
	Regular	9	2.9	1.2	
HDL (mmol/l)	irregular	23	3.6	1.7	0.4
	Regular	9	4.3	0.5	
LDL (mmol/l)	irregular	23	4.7	2.4	0.4
	Regular	9	4.2	1.3	

\*The mean difference is significant at the (0.05) level.

Table 15. Association between infertility hormones& lipid profile with BMI in patients.

BMI		N	Mean	Std. Deviation	P.value
FSH (ng/dl)	Normal	33	5.3	3.2	0.4
	overweight	7	3.9	1.7	
LH (ng/dl)	Normal	33	7.8	2.7	0.4
	overweight	7	6.3	3.0	
PRL (ng/dl)	Normal	33	31.4	13.9	0.8
	overweight	7	30.0	5.9	
CHO (mmol/l)	Normal	33	5.6	1.3	0.2
	overweight	7	4.6	1.3	
TG (mmol/l)	Normal	33	2.9	1.9	0.1
	overweight	7	5.3	2.4	
HDL (mmol/l)	Normal	33	3.8	1.8	0.3
	overweight	7	4.9	0.4	
LDL (mmol/l)	Normal	33	5.0	2.1	0.2
	overweight	7	4.1	1.6	

\*The mean difference is significant at the (0.05) level.

study provides several notable insights into the clinical features, hormonal profiles, lipid levels, and cystic characteristics associated with PCOS in this population.

Menstrual abnormalities were significantly more prevalent in the case group (75%) compared to controls (67.5%), consistent with the established role of PCOS in causing menstrual dysfunction. A substantial portion of the case group (60%) also reported a positive family history, reinforcing the genetic and familial contribution to PCOS pathogenesis. Despite this, no statistically significant association was found between family history and cyst location ( $p = 0.5$ ), suggesting that cyst laterality may not be directly influenced by familial factors in this cohort.

In the case group, bilateral ovarian cysts were slightly more common (53%) than unilateral cysts (47%). However, there was no statistically significant association between cyst location and most clinical variables, including menstruation status ( $p = 0.2$ ), medical follow-up ( $p = 0.1$ ), and history of delivery ( $p = 0.5$ ). A significant finding was observed between cyst location and chronic disease status ( $p = 0.04$ ), with bilateral cysts being more associated with diabetes mellitus, indicating a potential link between more extensive ovarian involvement and metabolic disturbances.

The majority of participants in the case group had normal BMI (83%), with 17% classified as overweight. Despite this, lipid abnormalities were prominent. The case group demonstrated significantly elevated levels of total cholesterol, triglycerides (TG), HDL, and PRL compared to controls (all  $p < 0.01$ ), while LDL levels did not differ significantly. This suggests a PCOS-related dyslipidemia that may occur independently of BMI, aligning with evidence that PCOS is a metabolic disorder even in non-obese women.

When assessing the relationship between BMI and biochemical markers, no significant differences were observed, although overweight individuals tended to have higher TG levels (mean = 5.3 mmol/L) compared to those with normal BMI (mean = 2.9 mmol/L), suggesting a trend worth further exploration.

Among infertility-related hormones, prolactin (PRL) levels were significantly higher in the case group (mean = 31.2 ng/dL) compared to controls (mean = 14.7 ng/dL,  $p = 0.00$ ), reflecting

hyperprolactinemia's possible role in PCOS pathophysiology. However, FSH and LH levels, while slightly elevated in the case group, did not differ significantly.

Notably, LH levels were significantly associated with cyst location ( $p = 0.01$ ), with unilateral cysts showing higher LH levels, suggesting a potential link between hormonal regulation and the extent of ovarian involvement. However, other hormonal markers (FSH, PRL) and lipid profiles did not significantly vary with cyst laterality.

More than two-thirds of the case group had a history of delivery (68%), and half reported regular medical follow-up. These findings reflect varied reproductive outcomes among women with PCOS and indicate that access to care may not be uniform across the population.

### Conclusion.

This study highlights significant hormonal and lipid profile alterations in Sudanese women with PCOS, even in those with normal BMI. While cyst laterality does not appear to correlate strongly with most clinical parameters, its association with LH and chronic disease warrants further investigation. The high prevalence of menstrual irregularities, positive family history, and dyslipidemia emphasizes the need for early screening and multidisciplinary management of PCOS, particularly in regions where awareness and access to care may be limited.

### Acknowledgment.

We would like to express our gratitude to all participants involved in the study.

### Funding.

This study has not received any external funding.

### Conflict of interest.

The authors declare that there is no conflict of interest.

### REFERENCES

- Ding D.C, Chen W, Wang J.H, et al. Association between polycystic ovarian syndrome and endometrial, ovarian, and breast cancer: A population-based cohort study in Taiwan. *Medicine*. 2018;97:e12608.

2. Abdullah AA, Ahmed M, Oladokun A. Prevalence of infertility in Sudan: A systematic review and meta-analysis. *Qatar Medical Journal*. 2021;3:47.
3. Franks S. Polycystic ovary syndrome. *N Engl J Med*. 1995;333:853-861.
4. Rotterdam ESHRE / ASRM – Sponsored PCOS Consensus workshop Group. Revised consensus on diagnostic criteria and long – term health risks related to polycystic ovary syndrome. *Fertile steril*. 2004;81:19-25.
5. Teede H.J, Tay C.T, Laven J.J, et al. Recommendations from the 2023 international evidence-based guideline for the assessment and management of polycystic ovary syndrome. *European journal of endocrinology*. 2023;189:G43-G64.
6. Ovalle F, Azziz R. Insulin resistance, polycystic ovary syndrome, and type 2 diabetes mellitus. *Fertility and sterility*. 2002;77:1095-1105.
7. Hazlehurst J.M, Singh P, Bhogal G, et al. How to manage weight loss in women with obesity and PCOS seeking fertility?. *Clinical endocrinology*. 2022;97:208-216.
8. Yin X, Ji Y, Chan C.L.W, et al. The mental health of women with polycystic ovary syndrome: a systematic review and meta-analysis. *Archives of women's mental health*. 2021;24:11-27.