

GEORGIAN MEDICAL NEWS

ISSN 1512-0112

NO 6 (375) Июнь 2026

ТБИЛИСИ - NEW YORK



ЕЖЕМЕСЯЧНЫЙ НАУЧНЫЙ ЖУРНАЛ

Медицинские новости Грузии
საქართველოს სამედიცინო სიახლენი

GEORGIAN MEDICAL NEWS

Monthly Georgia-US joint scientific journal published both in electronic and paper formats of the Agency of Medical Information of the Georgian Association of Business Press.
Published since 1994. Distributed in NIS, EU and USA.

GMN: Georgian Medical News is peer-reviewed, published monthly journal committed to promoting the science and art of medicine and the betterment of public health, published by the GMN Editorial Board since 1994. GMN carries original scientific articles on medicine, biology and pharmacy, which are of experimental, theoretical and practical character; publishes original research, reviews, commentaries, editorials, essays, medical news, and correspondence in English and Russian.

GMN is indexed in MEDLINE, SCOPUS, PubMed and VINITI Russian Academy of Sciences. The full text content is available through EBSCO databases.

GMN: Медицинские новости Грузии - ежемесячный рецензируемый научный журнал, издаётся Редакционной коллегией с 1994 года на русском и английском языках в целях поддержки медицинской науки и улучшения здравоохранения. В журнале публикуются оригинальные научные статьи в области медицины, биологии и фармации, статьи обзорного характера, научные сообщения, новости медицины и здравоохранения. Журнал индексируется в MEDLINE, отражён в базе данных SCOPUS, PubMed и ВИНТИ РАН. Полнотекстовые статьи журнала доступны через БД EBSCO.

GMN: Georgian Medical News – საქართველოს სამედიცინო სიახლენი – არის ყოველთვიური სამეცნიერო სამედიცინო რეცენზირებადი ჟურნალი, გამოიცემა 1994 წლიდან, წარმოადგენს სარედაქციო კოლეგიისა და აშშ-ის მეცნიერების, განათლების, ინდუსტრიის, ხელოვნებისა და ბუნებისმეტყველების საერთაშორისო აკადემიის ერთობლივ გამოცემას. GMN-ში რუსულ და ინგლისურ ენებზე ქვეყნდება ექსპერიმენტული, თეორიული და პრაქტიკული ხასიათის ორიგინალური სამეცნიერო სტატიები მედიცინის, ბიოლოგიისა და ფარმაციის სფეროში, მიმოხილვითი ხასიათის სტატიები.

ჟურნალი ინდექსირებულია MEDLINE-ის საერთაშორისო სისტემაში, ასახულია SCOPUS-ის, PubMed-ის და ВИНТИ РАН-ის მონაცემთა ბაზებში. სტატიების სრული ტექსტი ხელმისაწვდომია EBSCO-ს მონაცემთა ბაზებიდან.

WEBSITE

www.geomednews.com

К СВЕДЕНИЮ АВТОРОВ!

При направлении статьи в редакцию необходимо соблюдать следующие правила:

1. Статья должна быть представлена в двух экземплярах, на русском или английском языках, напечатанная через **полтора интервала на одной стороне стандартного листа с шириной левого поля в три сантиметра**. Используемый компьютерный шрифт для текста на русском и английском языках - **Times New Roman (Кириллица)**, для текста на грузинском языке следует использовать **AcadNusx**. Размер шрифта - **12**. К рукописи, напечатанной на компьютере, должен быть приложен CD со статьей.

2. Размер статьи должен быть не менее десяти и не более двадцати страниц машинописи, включая указатель литературы и резюме на английском, русском и грузинском языках.

3. В статье должны быть освещены актуальность данного материала, методы и результаты исследования и их обсуждение.

При представлении в печать научных экспериментальных работ авторы должны указывать вид и количество экспериментальных животных, применявшиеся методы обезболивания и усыпления (в ходе острых опытов).

4. К статье должны быть приложены краткое (на полстраницы) резюме на английском, русском и грузинском языках (включающее следующие разделы: цель исследования, материал и методы, результаты и заключение) и список ключевых слов (key words).

5. Таблицы необходимо представлять в печатной форме. Фотокопии не принимаются. **Все цифровые, итоговые и процентные данные в таблицах должны соответствовать таковым в тексте статьи.** Таблицы и графики должны быть озаглавлены.

6. Фотографии должны быть контрастными, фотокопии с рентгенограмм - в позитивном изображении. Рисунки, чертежи и диаграммы следует озаглавить, пронумеровать и вставить в соответствующее место текста **в tiff формате**.

В подписях к микрофотографиям следует указывать степень увеличения через окуляр или объектив и метод окраски или импрегнации срезов.

7. Фамилии отечественных авторов приводятся в оригинальной транскрипции.

8. При оформлении и направлении статей в журнал МНГ просим авторов соблюдать правила, изложенные в «Единых требованиях к рукописям, представляемым в биомедицинские журналы», принятых Международным комитетом редакторов медицинских журналов - <http://www.spinesurgery.ru/files/publish.pdf> и http://www.nlm.nih.gov/bsd/uniform_requirements.html. В конце каждой оригинальной статьи приводится библиографический список. В список литературы включаются все материалы, на которые имеются ссылки в тексте. Список составляется в алфавитном порядке и нумеруется. Литературный источник приводится на языке оригинала. В списке литературы сначала приводятся работы, написанные знаками грузинского алфавита, затем кириллицей и латиницей. Ссылки на цитируемые работы в тексте статьи даются в квадратных скобках в виде номера, соответствующего номеру данной работы в списке литературы. Большинство цитированных источников должны быть за последние 5-7 лет.

9. Для получения права на публикацию статья должна иметь от руководителя работы или учреждения визу и сопроводительное отношение, написанные или напечатанные на бланке и заверенные подписью и печатью.

10. В конце статьи должны быть подписи всех авторов, полностью приведены их фамилии, имена и отчества, указаны служебный и домашний номера телефонов и адреса или иные координаты. Количество авторов (соавторов) не должно превышать пяти человек.

11. Редакция оставляет за собой право сокращать и исправлять статьи. Корректуре авторам не высылаются, вся работа и сверка проводится по авторскому оригиналу.

12. Недопустимо направление в редакцию работ, представленных к печати в иных издательствах или опубликованных в других изданиях.

При нарушении указанных правил статьи не рассматриваются.

REQUIREMENTS

Please note, materials submitted to the Editorial Office Staff are supposed to meet the following requirements:

1. Articles must be provided with a double copy, in English or Russian languages and typed or computer-printed on a single side of standard typing paper, with the left margin of 3 centimeters width, and 1.5 spacing between the lines, typeface - **Times New Roman (Cyrillic)**, print size - 12 (referring to Georgian and Russian materials). With computer-printed texts please enclose a CD carrying the same file titled with Latin symbols.

2. Size of the article, including index and resume in English, Russian and Georgian languages must be at least 10 pages and not exceed the limit of 20 pages of typed or computer-printed text.

3. Submitted material must include a coverage of a topical subject, research methods, results, and review.

Authors of the scientific-research works must indicate the number of experimental biological species drawn in, list the employed methods of anesthetization and soporific means used during acute tests.

4. Articles must have a short (half page) abstract in English, Russian and Georgian (including the following sections: aim of study, material and methods, results and conclusions) and a list of key words.

5. Tables must be presented in an original typed or computer-printed form, instead of a photocopied version. **Numbers, totals, percentile data on the tables must coincide with those in the texts of the articles.** Tables and graphs must be headed.

6. Photographs are required to be contrasted and must be submitted with doubles. Please number each photograph with a pencil on its back, indicate author's name, title of the article (short version), and mark out its top and bottom parts. Drawings must be accurate, drafts and diagrams drawn in Indian ink (or black ink). Photocopies of the X-ray photographs must be presented in a positive image in **tiff format**.

Accurately numbered subtitles for each illustration must be listed on a separate sheet of paper. In the subtitles for the microphotographs please indicate the ocular and objective lens magnification power, method of coloring or impregnation of the microscopic sections (preparations).

7. Please indicate last names, first and middle initials of the native authors, present names and initials of the foreign authors in the transcription of the original language, enclose in parenthesis corresponding number under which the author is listed in the reference materials.

8. Please follow guidance offered to authors by The International Committee of Medical Journal Editors guidance in its Uniform Requirements for Manuscripts Submitted to Biomedical Journals publication available online at: http://www.nlm.nih.gov/bsd/uniform_requirements.html
http://www.icmje.org/urm_full.pdf

In GMN style for each work cited in the text, a bibliographic reference is given, and this is located at the end of the article under the title "References". All references cited in the text must be listed. The list of references should be arranged alphabetically and then numbered. References are numbered in the text [numbers in square brackets] and in the reference list and numbers are repeated throughout the text as needed. The bibliographic description is given in the language of publication (citations in Georgian script are followed by Cyrillic and Latin).

9. To obtain the rights of publication articles must be accompanied by a visa from the project instructor or the establishment, where the work has been performed, and a reference letter, both written or typed on a special signed form, certified by a stamp or a seal.

10. Articles must be signed by all of the authors at the end, and they must be provided with a list of full names, office and home phone numbers and addresses or other non-office locations where the authors could be reached. The number of the authors (co-authors) must not exceed the limit of 5 people.

11. Editorial Staff reserves the rights to cut down in size and correct the articles. Proof-sheets are not sent out to the authors. The entire editorial and collation work is performed according to the author's original text.

12. Sending in the works that have already been assigned to the press by other Editorial Staffs or have been printed by other publishers is not permissible.

**Articles that Fail to Meet the Aforementioned
Requirements are not Assigned to be Reviewed.**

ავტორთა საყურადღებო!

რედაქციაში სტატიის წარმოდგენისას საჭიროა დავიცვათ შემდეგი წესები:

1. სტატია უნდა წარმოადგინოთ 2 ცალად, რუსულ ან ინგლისურ ენებზე, დაბეჭდილი სტანდარტული ფურცლის 1 გვერდზე, 3 სმ სიგანის მარცხენა ველისა და სტრიქონებს შორის 1,5 ინტერვალის დაცვით. გამოყენებული კომპიუტერული შრიფტი რუსულ და ინგლისურენოვან ტექსტებში - **Times New Roman (Кириллица)**, ხოლო ქართულენოვან ტექსტში საჭიროა გამოვიყენოთ **AcadNusx**. შრიფტის ზომა – 12. სტატიას თან უნდა ახლდეს CD სტატიით.

2. სტატიის მოცულობა არ უნდა შეადგენდეს 10 გვერდზე ნაკლებს და 20 გვერდზე მეტს ლიტერატურის სიის და რეზიუმეების (ინგლისურ, რუსულ და ქართულ ენებზე) ჩათვლით.

3. სტატიაში საჭიროა გაშუქდეს: საკითხის აქტუალობა; კვლევის მიზანი; საკვლევი მასალა და გამოყენებული მეთოდები; მიღებული შედეგები და მათი განსჯა. ექსპერიმენტული ხასიათის სტატიების წარმოდგენისას ავტორებმა უნდა მიუთითონ საექსპერიმენტო ცხოველების სახეობა და რაოდენობა; გაუტკივარებისა და დაძინების მეთოდები (მწვავე ცდების პირობებში).

4. სტატიას თან უნდა ახლდეს რეზიუმე ინგლისურ, რუსულ და ქართულ ენებზე არანაკლებ ნახევარი გვერდის მოცულობისა (სათაურის, ავტორების, დაწესებულების მითითებით და უნდა შეიცავდეს შემდეგ განყოფილებებს: მიზანი, მასალა და მეთოდები, შედეგები და დასკვნები; ტექსტუალური ნაწილი არ უნდა იყოს 15 სტრიქონზე ნაკლები) და საკვანძო სიტყვების ჩამონათვალი (key words).

5. ცხრილები საჭიროა წარმოადგინოთ ნაბეჭდი სახით. ყველა ციფრული, შემავსებელი და პროცენტული მონაცემები უნდა შეესაბამებოდეს ტექსტში მოყვანილს.

6. ფოტოსურათები უნდა იყოს კონტრასტული; სურათები, ნახაზები, დიაგრამები - დასათაურებული, დანომრილი და სათანადო ადგილას ჩასმული. რენტგენოგრაფიის ფოტოსურათები წარმოადგინეთ პოზიტიური გამოსახულებით **tiff** ფორმატში. მიკროფოტოსურათების წარწერებში საჭიროა მიუთითოთ ოკულარის ან ობიექტივის საშუალებით გადიდების ხარისხი, ანათალების შედეგების ან იმპრეგნაციის მეთოდი და აღნიშნოთ სურათის ზედა და ქვედა ნაწილები.

7. სამამულო ავტორების გვარები სტატიაში აღინიშნება ინიციალების თანდართვით, უცხოურისა – უცხოური ტრანსკრიპციით.

8. სტატიას თან უნდა ახლდეს ავტორის მიერ გამოყენებული სამამულო და უცხოური შრომების ბიბლიოგრაფიული სია (ბოლო 5-8 წლის სიღრმით). ანბანური წყობით წარმოდგენილ ბიბლიოგრაფიულ სიაში მიუთითეთ ჯერ სამამულო, შემდეგ უცხოელი ავტორები (გვარი, ინიციალები, სტატიის სათაური, ჟურნალის დასახელება, გამოცემის ადგილი, წელი, ჟურნალის №, პირველი და ბოლო გვერდები). მონოგრაფიის შემთხვევაში მიუთითეთ გამოცემის წელი, ადგილი და გვერდების საერთო რაოდენობა. ტექსტში კვადრატულ ფხიხლებში უნდა მიუთითოთ ავტორის შესაბამისი N ლიტერატურის სიის მიხედვით. მიზანშეწონილია, რომ ციტირებული წყაროების უმეტესი ნაწილი იყოს 5-6 წლის სიღრმის.

9. სტატიას თან უნდა ახლდეს: ა) დაწესებულების ან სამეცნიერო ხელმძღვანელის წარდგინება, დამოწმებული ხელმოწერითა და ბეჭდით; ბ) დარგის სპეციალისტის დამოწმებული რეცენზია, რომელშიც მითითებული იქნება საკითხის აქტუალობა, მასალის საკმაობა, მეთოდის სანდოობა, შედეგების სამეცნიერო-პრაქტიკული მნიშვნელობა.

10. სტატიის ბოლოს საჭიროა ყველა ავტორის ხელმოწერა, რომელთა რაოდენობა არ უნდა აღემატებოდეს 5-ს.

11. რედაქცია იტოვებს უფლებას შეასწოროს სტატია. ტექსტზე მუშაობა და შეჯერება ხდება საავტორო ორიგინალის მიხედვით.

12. დაუშვებელია რედაქციაში ისეთი სტატიის წარდგენა, რომელიც დასაბეჭდად წარდგენილი იყო სხვა რედაქციაში ან გამოქვეყნებული იყო სხვა გამოცემებში.

აღნიშნული წესების დარღვევის შემთხვევაში სტატიები არ განიხილება.

E. Tsitlidze, I. Rukhadze, I. Verulashvili, Derry Minyao NG. CEREBROVASCULAR EVENTS AND NEUROCOGNITIVE IMPAIRMENTS WITH ALTERATIONS IN POLYSOMNOGRAPHIC INDICATORS FOLLOWING CARDIAC SURGERY.....	6-15
Saleh Alsuwaydani. TRENDS IN GENERAL SURGERY PROCEDURE TYPES AND THEIR OUTCOMES FOR A NEWLY COMMISSIONED HOSPITAL.....	16-20
Maltsev Dmytro. BEYOND THE GENETIC CODE: SYSTEMIC REGULATORY MELTDOWN AS A FRAMEWORK FOR PRIMARY EPIGENETIC DISEASES.....	21-42
David Malidze, Lasha Dolidze, Natia Jojua, Tinatin Gognadze, Ana Jabauri, Eka Kokhraidze, Larisa Melia, Tamar Chanturia, Tamar Chkoidze, Stephen Charles Jones. STANDARDIZED PATIENT-BASED TRAINING AND CLINICAL DIAGNOSTIC COMPETENCE AMONG MEDICAL STUDENTS: A RANDOMIZED COMPARATIVE STUDY.....	43-50
Ghukasyan Norayr, Melik-Andreasyan Marine, Sahakyan Lusine. HODGKIN LYMPHOMA DIAGNOSED DURING THE SECOND TRIMESTER OF PREGNANCY AND TREATED WITH ABVD CHEMOTHERAPY WITH FAVORABLE MATERNAL AND NEONATAL OUTCOMES: A CASE REPORT.....	51-54
Tchatchia Gr, Chelishvili I, Chutkerashvili G, Inauri N, Mermanishvili T. THE EFFECTIVENESS OF INTRAOPERATIVE ULTRASOUND IN GLIAL TUMOR SURGERY.....	55-62
Rakhmetullina A.K, Alimbayeva A.R, Orekhov A.Yu, Dairbekov Ye.Ye, Rakhmetullin A.B. COGNITIVE IMPAIRMENT AFTER CORONARY ARTERY BYPASS GRAFTING: THE ROLE OF MEDICAL REHABILITATION — A LITERATURE REVIEW.....	63-72
N.G. Alibeyli, N.M. Amiraliyev, H.K. Muradov, N.V. Kerimova, K.N. Amiraliyev, A.T. Iskenderova, E.N. Mekhtiyeva, S.D. Huseynova. GIANT CUTANEOUS HORN OF THE CHEEK: A CLINICAL CASE REPORT.....	73-76
Samara S. Yonus, Shamil H. Othman, Salah Yousif, Marwan Merkhani. NEUROPROTECTIVE FITNESS OF LOSARTAN AGAINST DOXORUBICIN INDUCED NEUROTOXICITY IN WHITE ALBINO RATS.....	77-83
Kurbanov Anvar Alamovich, Abdullaeva Yayra Davronovna, Matluba Badritdinova, Abdullaeva Vasila Karimbekovna, Khalimova Fariza Tursunbaevna, Marwan Ismail. DYNAMIC BIOMARKERS FOR PREDICTING POST-TRAUMATIC EPILEPSY AFTER TRAUMATIC BRAIN INJURY.....	84-89
J. Orozco Pilco, C. Calderón Cabezas, L. Santander Samaniego, G. Damián Sinchiguano A COMPARATIVE ANALYSIS OF TRENDS IN CERVICAL CANCER MORTALITY AMONG MESTIZO AND INDIGENOUS WOMEN IN ECUADOR, 2010–2024.....	90-95
Ling-Ling Zhou, Lian-Ping He. DUAL-SOFTWARE INTEGRATION IN MEDICAL STATISTICS TEACHING: A CASE-BASED APPROACH TO F-DISTRIBUTION WITH DOUBAO AI AND R.....	96-99
Essmat J. Jamel, Ayoub Ali Hussein Al-bayti, Alaa E. Jamal, Asmaa E. Jamal. ASSESSMENT OF VITEX AGNUS-COSTES PLANT EXTRACT ON THE HEART OF MICE EXPOSED TO OXIDATIVE STRESS.....	100-110
Nana Chikladze, Salome Kordzaia, Lika Svanadze, Tamar Rukhadze, Carlos Centeno. CRYOTHERAPY FOR THE PREVENTION OF TAXANE-INDUCED PERIPHERAL NEUROPATHY IN BREAST CANCER: A PROSPECTIVE MULTICENTER STUDY.....	111-117
Saipudinkyzy A, Yessirkepov Assilbek, Aitbayev Zhanabay, Laktionova M, Baimuratova M. PSYCHOMETRIC ASSESSMENT OF AN ADAPTED BILINGUAL (RUSSIAN–KAZAKH) VERSION OF THE UW-QOL V4.1 QUESTIONNAIRE AND ANALYSIS OF FUNCTIONAL REHABILITATION OUTCOMES IN PATIENTS OF THE REPUBLIC OF KAZAKHSTAN.....	118-127
Andriy Kyrylyuk, Mykola Rozhko, Mykola Kyrylyuk, Liubow Kyryliuk, Olena Rozhko. EFFECTIVENESS OF STANDARD AND INDIVIDUALIZED MARPE DEVICES IN ADULT PATIENTS BASED ON CONE BEAM COMPUTED TOMOGRAPHY DATA.....	128-137
Issametov Davran Rashitovich, Gantsev Kamil Shamilevich, Arybzhonov Dauranbek Tursunculovich, Osman Kostek, Kemelbekov Kanatzhon Saukhanbekovich, Balakhnin Pavel Vasilyevich, Saburov Alisher Radzhapbaevich, Saburova Saidokhon Alisherovna. SELECTIVE BRONCHIAL ARTERY CHEMOINFUSION IN NSCLC: CURRENT STATE OF REGIONAL THERAPY AND ITS ROLE IN MULTIMODAL APPROACHES.....	138-146
Rasha A.H. Alathary, Nawal Khintee Jabbar, Sundus K. Hamzah. ASSESSMENT OF SUBCLINICAL RENAL INSULT IN SHORT - AND LONG-TERM UNCONTROLLED HYPERTENSION USING INTEGRATED MOLECULAR INDEX OF NGAL AND KIM-1.....	147-152

Olena Kozeratska, Adelina Gutnitska, Oksana Vdovichenko, Vitalii Spivak, Alisa Kabesheva. MENTAL HEALTH IN THE MODERN WORLD: THE PROBLEM AND SOLUTIONS.....	153-160
Zurab S. Khabadze, Magomed-Ali A. Gasbanov, Nasrula R. Akhmedov, David A. Babakhanov, Khamzat A. Umarov, Nikita A. Dolzhikov, Eliso M. Kakabadze, Nadezhda A. Khachatryan, Veranika M. Slonova. MANAGEMENT OF CHRONIC APICAL PERIODONTITIS WITH FURCATION INVOLVEMENT USING HIGH-TEMPERATURE SODIUM HYPOCHLORITE IRRIGATION: A CASE REPORT.....	161-165
Hafiza S. Zhetpisbayeva, Karashash M. Askarova, Aigerim E. Tolendiyeva, Kalima T. Kupenova, Baglan O. Zharmagambetova, Nurzhamal D. Imambayeva, Zhaniya I. Zhakypova. CLINICAL CHARACTERISTICS AND RISK FACTORS FOR ADVERSE OUTCOMES IN ELDERLY PATIENTS WITH SEVERE COVID-19.....	166-174
Lavdie Leci, Berat Lenjani, Ilijana Muratovska. VOLUMETRIC AND RADIOGRAPHIC OUTCOMES OF BONE REGENERATION AFTER APICOECTOMY WITH PRF-BASED THERAPIES - A PRELIMINARY STUDY.....	175-182
María Evelyn Alviárez, Mario Alexander Aguilar Balseca, Ana Carolina González Romero, Betzabeth Thayin Guillen Armijo, Eva Iralic Quiñonez Ruiz. ANTIMICROBIAL SUSCEPTIBILITY PATTERNS OF <i>ESCHERICHIA COLI</i> AND <i>ENTEROBACTER SPP.</i> CAUSING COMMUNITY-ACQUIRED INFECTIONS IN MÉRIDA, VENEZUELA.....	183-188
Nino Totadze, Nino Adamia, Manana Kobakhidze, Irine Kekelidze, Julieta Kikvadze, Lasha Tchelidze. SKIN-TO-SKIN CONTACT IN NEONATAL CARE: CLINICAL SIGNIFICANCE, IMPLEMENTATION CHALLENGES, AND LONG-TERM OUTCOMES IN GEORGIA.....	189-198
L. Stepanyan, E. Asrian, G. Lalayan, N. Harutyunyan. ACADEMIC ENGAGEMENT AS A CONTEMPORARY CHALLENGE IN HIGHER EDUCATION: TOWARD A FUNCTIONAL INTEGRATIVE MODEL.....	199-206
Guranda Okropiridze, Arsen Gvenetadze, Revaz Sulukhia, Rusudan Gvenetadze, Diana Chanukvadze, Lela Iremadze, Ana Gvenetadze, Maka Gegechkori, Nino Kikvadze. IMPACT OF THE FIRST TRIMESTER PREECLAMPSIA SCREENING TEST ON THE OUTCOME OF PREGNANCY BETWEEN THE PATIENTS WITH OR WITHOUT THE DIAGNOSIS OF INFERTILITY.....	207-211
Yerlan A. Salmenbayev, Yessengazy O. Massalimov, Altai A. Dyussupov, Nazarbek B. Omarov, Assem D. Kazangapova. PREVENTION OF LOCAL POSTOPERATIVE COMPLICATIONS IN THE SURGICAL TREATMENT OF CHRONIC VENOUS INSUFFICIENCY OF THE LOWER EXTREMITIES (CEAP C3–C5).....	212-221
Aws S. Sheet, Mohammed KJ. Alnori. SAFETY OF TAMSULOSIN ALONE OR IN COMBINATION WITH 5 α -REDUCTASE INHIBITORS ON LIPID PROFILE, RENAL FUNCTION, AND LIVER PARAMETERS.....	222-229
Zeyad Hussain Althumali, Jumana Hamed Khouja, Reem F Alnemari, Moteab Sayer Alotaybi, Fawziah Roublah, Asma M. Alahmari, Abdullah H. Jabbar, Adeeb A Almohammadi, Hatem mazen Alsolami. ASSESSING THE EFFECTIVENESS OF LIFESTYLE CLINIC INTERVENTIONS ON CARDIOMETABOLIC RISK FACTORS: A RETROSPECTIVESTUDY.....	230-237
Natia Nazghaidze, Rusudan Agladze, Zviad Kereselidze, Nika Kuridze, Sophio Ungiadze, Shalva Petriashvili. EFFICACY OF PERINDOPRIL IN PATIENTS WITH HEART FAILURE AND REDUCED EJECTION FRACTION (HFrEF)	238-247

GIANT CUTANEOUS HORN OF THE CHEEK: A CLINICAL CASE REPORT

N.G. Alibeyli, N.M. Amiraliyev, H.K. Muradov, N.V. Kerimova, K.N. Amiraliyev, A.T. Iskenderova, E.N. Mekhtiyeva, S.D. Huseynova.

Department of Oncology, Azerbaijan Medical University, Baku, Azerbaijan.

Abstract.

A cutaneous horn is a conical keratinous projection that may develop on various parts of the body, most commonly in sun-exposed areas. These lesions are of considerable clinical significance because they may be associated with premalignant or malignant conditions. Histologically, cutaneous horns consist of hyperkeratotic material and resemble animal horns; however, they lack the central bony core characteristic of true animal horns.

We present the case of a male patient with a giant cutaneous horn located on the skin of the right cheek. The lesion was managed by wide local excision followed by reconstruction of the resulting defect. Histopathological examination revealed an underlying malignant process, specifically squamous cell carcinoma.

The standard treatment for a cutaneous horn is complete surgical excision with adequate margins, followed by histopathological evaluation to determine the need for further management and appropriate therapeutic strategies.

Key words. Giant cutaneous horn, carcinoma, radical excision, reconstruction.

Introduction.

A cutaneous horn (*cornu cutaneum*) may represent a benign, premalignant, or malignant skin lesion and is most commonly observed in areas chronically exposed to ultraviolet radiation. Although cutaneous horns can occur on any part of the body, approximately 30% of cases are located on the face and scalp. Surgical excision remains the treatment of choice, particularly for facial lesions, where the likelihood of underlying malignancy is considerably higher. Histopathological examination of the base of the horn is essential to exclude malignant transformation and to establish the definitive diagnosis [1-4].

Squamous cell carcinoma presenting as a giant cutaneous horn of the cheek is an exceptionally rare condition. To the best of our knowledge, the present case represents one of the largest cutaneous horns of the cheek reported in the literature.

Case Report.

A 68-year-old man was admitted to our department with a painless, slowly enlarging lesion on the skin of the right cheek. The lesion had progressively evolved from a small mass into a giant cutaneous horn over a period of five years.

The patient had no previous history of skin malignancy. The lesion demonstrated a prolonged course characterized by continuous slow growth. Although surgical treatment had been recommended earlier, the patient declined the procedure and underwent topical treatment prescribed by a dermatologist, without clinical improvement.

On physical examination, a giant horn-like lesion measuring 8 × 6 cm was observed on a broad base. No regional

lymphadenopathy or bleeding was detected. During the last year, the lesion had become a significant cosmetic concern for the patient (Figure 1).

Routine laboratory investigations were within normal limits. Computed tomography (CT) revealed no evidence of invasion into the bones of the facial skeleton (Figure 2).

Under general endotracheal anesthesia, the patient underwent wide surgical excision of the tumor with adequate margins (Figure 3). The postoperative soft-tissue defect was reconstructed using a local musculocutaneous flap, providing satisfactory coverage and restoration of facial contour (Figures 4-6).

Postoperative Course.

No local complications were observed during the postoperative period. The wound healed uneventfully by primary intention (Figure 7). Histopathological examination of the excised specimen confirmed the diagnosis of squamous cell carcinoma (Figure 8).

Discussion.

A cutaneous horn is a clinical diagnosis referring to a conical projection arising above the surface of the skin. It resembles an animal horn and consists of compacted keratinized material [2]. Although cutaneous horns macroscopically resemble animal horns, they differ histologically. Animal horns are composed of a hyperkeratotic epidermis covering a dermal core with a centrally located, well-developed bony structure, whereas human cutaneous horns, including giant lesions, lack a central osseous core [5-11].

Cutaneous horns have been reported predominantly in Caucasian and Asian populations and are relatively uncommon among individuals of African descent [1]. More than 30% of cases occur in the head and neck region [2,12-16]. Other sun-exposed areas, including the chest, neck, shoulders, and dorsum of the hands, are also common sites for the development of cutaneous horns.

The principal clinical concern is not the horn itself, which consists of dead keratin, but rather the underlying pathological process at its base. The associated lesion may be benign, premalignant, or malignant. Benign conditions include seborrheic keratosis, viral warts, histiocytoma, epidermal nevus, and molluscum contagiosum. Premalignant lesions include actinic keratosis and Bowen's disease, while malignant lesions most commonly comprise squamous cell carcinoma and, less frequently, basal cell carcinoma or Kaposi's sarcoma [6].

Cutaneous horns exhibit considerable variation in size and morphology and may be cylindrical, conical, or curved in shape [2]. Previous reports have suggested that large conical horns, particularly giant cutaneous horns, are more likely to arise from an underlying malignant lesion [9]. Cutaneous horns occur most frequently in individuals older than 50 years and affect both



Figure 1. Giant cutaneous horn arising from the skin of the right cheek.



Figure 4. Intraoperative view demonstrating flap elevation and pedicle division.



Figure 2. Preoperative computed tomography (CT) imaging of the patient.



Figure 5. The pedicled flap was elevated and transposed to the defect through a subcutaneous tunnel created above the lower border of the mandible.



Figure 3. Postoperative defect of the right cheek after wide local excision of the tumor with a 1-cm margin. Preoperative design and marking of an island platysma musculocutaneous flap for defect reconstruction.



Figure 6. Reconstruction of the cheek defect with flap inset and fixation using interrupted sutures. Flap perfusion and viability were confirmed by the needle-prick test.



Figure 7. Four-month postoperative follow-up showing good cosmetic outcome and absence of local recurrence.




Figure 8. Histological section revealing squamous cell carcinoma associated with a cutaneous horn.

sexes; however, malignant underlying pathology is observed more commonly in men [17].

The largest study to date, conducted by Yu et al. and including 643 cases of cutaneous horns, demonstrated that 39% of lesions originated from malignant or premalignant epidermal conditions, whereas 61% were associated with benign underlying pathology [17].

Matneše et al. [9], in a histopathological analysis of 222 cases of cutaneous horn, reported that 59% of lesions were associated with premalignant or malignant pathology. Among the malignant lesions, squamous cell carcinoma (SCC) accounted for 94.0% of cases. Similarly, Pyne et al. [12] found that SCC was the most common histopathological diagnosis at the base of the horn, occurring in 34% of patients (56 of 163 cases).

Çorçu et al. [3] analyzed 11 cases of cutaneous horn and identified two cases of squamous cell carcinoma, one case of basal cell carcinoma, and eight cases with benign pathology.

Spira and Rabinovitz [14] concluded that cutaneous horns associated with malignant or premalignant lesions occur more frequently in patients with a history of other malignant or premalignant skin conditions. In their series of 38 patients, 44% were found to have associated malignant neoplasms.

Mir et al. [10] reported that in three of six cases of cutaneous horn, histopathological examination revealed an underlying squamous cell carcinoma.

Squamous cell carcinoma presenting as a giant cutaneous horn of the cheek, as observed in our patient, is an exceptionally rare condition. To the best of our knowledge, this case represents one of the largest cutaneous horns of the cheek reported in the literature. Even among the largest previously documented cases, such as the massive cheek lesion described by Wang et al. [16], the maximum diameter of the lesion's base was reported to be 6 cm. A distinctive feature of our case is the exceptionally large tumor base, which measured 8 cm in diameter, making it one of the largest cheek-based cutaneous horns reported to date.

Conclusion.

A cutaneous horn is a keratinous skin lesion of variable size that most commonly develops on sun-exposed areas of the body. Premalignant and malignant conditions are frequently

identified at its base, highlighting the importance of thorough diagnostic evaluation. Complete surgical excision followed by histopathological examination remains the recommended approach for establishing an accurate diagnosis, guiding further management, and achieving satisfactory cosmetic outcomes.

REFERENCES

1. Akan M, Yildirim S, Avci G, et al. Xeroderma pigmentosum with a giant cutaneous horn. *Annals Plastic Surgery*. 2001;46:665-666.
2. Bondeson J. Everard Home, John Hunter, and cutaneous horns: a historical review. *American Journal of Dermatopathology*. 2001;23:362-369.
3. Copcu E, Sivrioglu N, Culhaci N. Cutaneous horns: are these lesions as innocent as they seem to be? *World Journal of Surgical Oncology*. 2004;2:1-4.
4. Fernandes N, Sinha S, Lambert C, et al. Cutaneous horn: a potentially malignant entity. *Acta Dermato venerol Alpina Pannonica Adriatica*. 2009;18:189-193.
5. İro S, Maadane A, Slimani F. Facial cutaneous horn in three different conditions: a highlight of a malignancy risk. *International Journal of Surgery Case Reports*. 2021;83:1-4.
6. Kitagawa H, Mizuno M, Nakamura Y, et al. Cutaneous horn can be a clinical manifestation of underlying sebaceous carcinoma. *British Journal of Dermatology*. 2007;156:180-182.
7. Kneitz H, Motschenbacher S, Wobser M, et al. Photoletter to the editor: giant cutaneous horn associated with squamous cell carcinoma. *Journal Dermatology Case Report*. 2015;9:27-28.
8. Kumar S, Bijalwan P, Saini S. Carcinoma buccal mucosa underlying a giant cutaneous horn: a case report and review of the literature. *Case Reports in Oncological Medicine*. 2014;11:38-40.
9. Mantese S, Diogo P, Rocha A, et al. Cutaneous horn: a retrospective histopathological study of 222 cases. *Anais Brasileiros de Dermatologia*. 2010;85:157-163.
10. Michal M, Bisceglia M, Di Mattia A, et al. Gigantic cutaneous horns of the scalp: lesions with a gross similarity to the horns of animals: a report of four cases. *American Journal of Surgical Pathology*. 2002;26:789-794.

11. Mir M, Mahmud A, Mohd Y, et al. Cutaneous horn: a devil not only in appearance. *Plastic Surgery Case Studies*. 2016;2:51-52.
12. Pyne J, Sapkota D, Wong JC. Cutaneous horns: clues to invasive squamous cell carcinoma being present in the horn base. *Dermatology Practical Conceptual*. 2013;3:3-7.
13. Sathyanarayana S.A, Deutsch G.B, Edelman M, et al. Cutaneous horn: a malignant lesion: A brief review of the literature. *Dermatology Surgery*. 2012;38:285-287.
14. Spira J, Rabinovitz H. Cutaneous horn present for two month. *Dermatology Online J*. 2000;6:11.
15. Vanik S, Mehta S, Shah R, et al. Giant cutaneous horn: is it a tip of an iceberg? two case reports and a review of literature. *Indian Journal Otolaryngol Head and Neck Surgery*. 2021;74:5432-5435.
16. Wang Y, Guo M, Liu B, et al. A giant cutaneous horn on the cheek. *Journal of Plastic Reconstructive & Aesthetic Surgery*. 2013;66:871-872.
17. Yu R, Pryce D, Macfarlane A, et al. A histopathological study of 643 cutaneous horns. *British Journal of Dermatology*. 1991;124:449-452.